

Vitamin D deficiency and its associating factors: A cross-sectional analytic study

Dr. Munmun Yadav¹, Dr. Sangeeta Kamra², Dr. Antarlina Ray³, Dr. Mahendra Kumar Verma^{4§}, Dr. Kusum Gaur⁵

^{1,3}DNB student, Department of Gynecology and Obstetrics, JLN Hospital and Research Centre Bhilai Steel Plant, Bhilai, Chhattisgarh, India

²Medical Senior Deputy Director, Department of Gynecology and Obstetrics, JLN Hospital and Research Centre Bhilai Steel Plant, Bhilai, Chhattisgarh, India

⁴Post graduate residents, Department of Community Medicine, SMS Medical College, Jaipur (Rajasthan) India

⁵Professor, Department of Community Medicine, SMS Medical College, Jaipur (Rajasthan) India

[§]Corresponding author's Email: verma.udr@gmail.com

Abstract— Vitamin D deficiency is widely prevalent throughout the world. Pregnant women, neonates and infants form most vulnerable groups for vitamin D deficiency. Hypovitaminosis vitamin D in pregnancy has been reported to cause various fetomaternal effect. So this study was conducted to find out proportion of vitamin D deficiency among pregnant women and to assess the effects and its associating factors. For this purpose 120 Pregnant women on their first visit to hospital irrespective of gestational age were taken. Apart from routine obstetrical investigation, serum vitamin D (total) level was estimated. All results were recorded and analyzed statically. It was observed that 84.1% were found to be vitamin D deficient. Mean age of vitamin D deficient group and non deficient group were 28.31 ± 3.86 and 26.37 ± 2.83 years respectively. Religion and parity of women was not found to be associated with vitamin D deficiency whereas age, type of diet and use of sunscreen was found to be associated. Vitamin D deficiency was found significantly more in elderly, vegetarians and users of sunscreen than their counterparts. Regarding maternal and neonatal outcomes, development of preeclampsia, LSCS as mode of delivery, delivery of pre term and LBW babies were having controversial findings. So it suggest much more researches in this regards.

Keywords: Feto-Maternal Outcome, Hypovitaminosis D, Maternal Blood Vitamin D, Vitamin D Status in Pregnancy.

I. INTRODUCTION

Vitamin D, is a lipid-soluble vitamin and prohormone which is known to play an important role in bone metabolism through regulation of calcium and phosphate homeostasis¹. Vitamin D traditionally known as “Anti- ricketic or sunshine vitamin”. It is a unique nutrient because it can be synthesized endogenously (skin) and it function as a hormone²

Vitamin D deficiency is prevalent worldwide and its unrecognized epidemic is common in all age groups ranging from 46-90%³⁻⁶. Various Indian studies have reported vitamin D deficiency in all age groups and involving both sexes. It's prevalence ranging between 40-93%.^{7,8} During pregnancy high prevalence of vitamin D deficiency has been reported from various region of world, ranging from 45 - 100%.⁴⁻⁶ and in Indian population it has been reported to the tune of 42-93%.^{9,10}

Studies regarding prevalence of vitamin D deficiency in antenatal women in India are mainly from Northern and southern India^{9,10} and very a few from central Part of India, thus this study has been undertaken in order to know the burden of vitamin D deficiency in antenatal women in ANCs of JLN Hospital and Research Centre Bhilai Steel Plant, Bhilai, Chhattisgarh, India.

II. METHODOLOGY

This hospital based observational study was conducted in the Department of Obstetrics and Gynaecology in collaboration with the department of nuclear medicine of Jawahar Lal Nehru Hospital and Research Centre, Bhilai, Chhattisgarh, India, from October 2015 to November 2016. Proper approval from the hospital ethical committee was taken for the study.

Antenatal women on their first visit to hospital, irrespective of period of gestational age were included in this study. ANCs on vitamin D supplementation in early pregnancy before attending this hospital and who had any disease like hypertension & diabetes mellitus were excluded from the study. ANCs who were not willing to participate in this study also excluded from study. Finally 120 ANCs were selected for this study.

Sample size was calculated 104 subjects at 95% confidence limit and 12% relative allowable error assuming as follows vitamin D deficiency in pregnant women 72.1%. So for this study 120 ANCs were included. study.

After taking written informed consent and general information, routine obstetrical investigation, serum vitamin D level estimation was done by fully automated Chemiluminescent immunoassay and Radioimmunoassay (RIA) of each of the subject. They were divide into following category as per Vitamin D status:-

Reference range:- recently revised institute of medicine's (IOM) criteria 2010 ¹¹

S. No.	Vitamin D Status	Vitamin D3 Level (ng/ml)
1	Vitamin D sufficiency	≥30
2	Vitamin D insufficiency	21-29
3	Vitamin D deficiency	20 or <20

For the study purpose all antenatal women will be divided in to two groups:

1. Group 1: Those who were having Vitamin D 30 and more than 30 i.e. **Vitamin D sufficient group**
2. Group 2: Those who were having Vitamin D less than 30 i.e. **Vitamin D deficient group**

Statistical data analysis: Data thus collected were entered in Microsoft Excel Sheet on same day so as to minimize data entry bias if any. Analyses were done by using SPSS version 24 (trial version). The results were presented as percentages and association was assessed by Chi-square test. P-value of less than 0.05 was taken as significant.

III. RESULTS

Majority 49(40.83%) of ANCs first visit was in 28-36 weeks of gestational age. Minimum 6(5%) patient's first visit was in first trimester (<12 week). Out of these 120 ANCs, 101 cases (84.1%) were found vitamin D deficient. (Figure 1)

The mean age of vitamin D deficient and sufficient group was found 28.31±3.86 and 26.37±2.83 years respectively. This difference in mean age was with significant difference (p<0.05). (Table 1)

Maximum number (114 (95%) of ANCs were Hindu. Out of 101 vitamin D deficient ANCs, 97 (96.03%) were Hindus whereas out of 19 vitamin D sufficient ANCs only 17 (89.47%) were Hindus. On

further analysis religion of ANC was not found to be associated with vitamin D status ($p > 0.05$). (Table 1)

Maximum number 81(67.5%) ANCs were vegetarian. Out of 101 vitamin D deficient ANCs, 75 (74.26%) were vegetarian whereas out of 19 vitamin D sufficient ANCs only 6 (31.58%) were vegetarians. This distribution of ANCs with vitamin D deficiency as per the type of diet was found statistically significant ($p < 0.001$) i.e. proportion of vitamin D deficient women were found significantly more ($p < 0.001$) in vegetarians. (Table 1)

Majority of ANCs, 84(70%) were belonged to primipara only 36 (30%) were multipara. Out of 101 vitamin D deficient ANCs, 71 (70.3%) were primipara whereas out of 19 vitamin D sufficient ANCs only 13 (68.42%) were primipara. On further analysis parity of ANC was not found to be associated with vitamin D status ($p > 0.05$). (Table 1)

Table 1
Comparison of Characteristics of Vitamin D Deficient and Vitamin D Sufficient Study subjects

Characteristics		Vitamin D Deficient (N=101)	Vitamin D Sufficient (N=19)	Significance
Age	(Mean ± SD) Years	28.31±3.86	26.37±2.83	P=0.039 S
Religion	Hindu	97	17	P=0.528 NS
	Muslim	4	2	
Type of Diet	Vegetarian	75	6	P<0.001 S
	Non vegetarian	26	13	
Parity	Primipara	71	13	P=0.913 NS
	Multipara	30	6	

Among these 120 ANCs, 69 (57.75%) were using sunscreen lotion/cream. Out of 101 vitamin D deficient ANCs, 65 (64.36%) were using sunscreen whereas out of 19 vitamin D sufficient ANCs only 4 (21.05%) were using sunscreen. On further analysis, Vitamin D deficiency was found significantly more in ANCs using sunscreen lotions ($p < 0.05$). (Figure 2)

Figure 1

Distribution of ANCs as per Vitamin D Status





Figure 2

Association of Vitamin D Status with Sunscreen lotion



Out of 120 ANCs, 33(27.5%) ANCs developed preeclampsia and rest 87 (72.5%) ANCs were normotensive. Out of 101 vitamin D deficient ANCs, 26 (25.74%) were with preeclampsia whereas out of 19 vitamin D sufficient ANCs only 7 (36.84%) were with preeclampsia. This distribution of preeclampsics in vitamin D deficient and vitamin d sufficient group was without any significant difference. ($p>0.05$). (Table 2)

In this study of 120 ANCs, only 2 (1.6%) ANCs developed GDM out of which one was vitamin D sufficient and another one was vitamin D deficient as number of cases who developed GDM was only 2 and majority (98.01%) of ANCs were euglycemics. This distribution of ANCs with GDM in vitamin D deficient and vitamin D sufficient group was without any significant difference. ($p>0.05$). (Table 2)

Out of 120 ANCs, 30 (29.7%) ANCs delivered by LSCS. Out of 101 vitamin D deficient ANCs, 23 (22.77%) had LSCS whereas out of 19 vitamin D sufficient ANCs only 7 (36.84%) had LSCS. This distribution was with significant difference ($p<0.05$). Surprisingly LSCS were found significantly more in ANCs with vitamin D sufficiency. (Table 2)

Table 2

Comparison of Maternal Complication in Vitamin D Deficient and Vitamin D Sufficient Study subjects

Characteristics		Vitamin D Deficient (N=101)	Vitamin D Sufficient (N=19)	Significance
Preeclampsia	Yes	26	7	P=0.475 NS
	No	75	12	
GDM	Yes	1	1	P=0.720 NS
	No	100	18	
LSCS	Yes	23	7	P=0.039 S
	No	78	6	

In the study of 120 ANCs 108 (90%) ANCs delivered at term and 12 (10%) ANCs gave birth to preterm baby. Out of 101 vitamin D deficient ANCs, 10 (9.9%) delivered preterm babies whereas out of 19 vitamin D sufficient ANCs only 2 (10.52%) delivered preterm babies. This difference in proportion of preterm babies born by vitamin D deficient ANCs and vitamin D sufficient ANCs was found not significant ($P>0.05$) (Table 3)

Out of 120 ANCs 35 (29.1%) ANCs delivered low birth weight babies. Out of 101 vitamin D deficient ANCs, 27 (26.73%) delivered LBW babies whereas out of 19 vitamin D sufficient ANCs only 8 (42.1%) delivered LBW babies. This difference in proportion of LBW babies born by vitamin D deficient ANCs and vitamin D sufficient ANCs was found not significant ($P>0.05$) (Table 3)

Table 3

Comparison of Neonatal Complication in Vitamin D Deficient and Vitamin D Sufficient Study subjects

Characteristics		Vitamin D Deficient (N=101)	Vitamin D Sufficient (N=19)	Significance
Preterm	Yes	10	2	P=0.739 NS
	No	91	17	
LBW	Yes	27	8	P=0.281 NS
	No	74	11	

IV. DISCUSSION

In this present study proportion of vitamin D deficiency was found 84.1%. Almost similarly observations were made by Jani R et al (2004),⁸ Sachan A et al (2005),⁹ Dasgupta et al (2011)¹⁰ and Rooplekha Chauhan et al (2015)¹² who reported 100%, 84.3%, 42% and 72.1% respectively.

Although religion and parity of ANC was not found to be associated with vitamin D deficiency in this present study but age & type of diet were found the associating factors. Vitamin D deficiency was found significantly higher in elder age group and vegetarians than their counterparts.

In present study, mean age of vitamin D deficient and sufficient group was found 28.31 ± 3.86 and 26.37 ± 2.83 years respectively, which was with significant difference. Johnson D et al (2015)¹³ conducted study on 310 pregnant women and found mean age 24.3 (range 18-40 year) of vitamin D deficient ANCs which was little lesser the present study.

Regarding type of diet observations of Crowel et al¹⁴ were well in resonance with present study who found that plasma 25(OH)D concentration were lower in vegetarian than meat and fish eater as in this present study also observed that vitamin D deficient ANCs were significantly higher in vegetarians.

In present study, although proportion of vitamin D deficient ANCs were little higher in primipara than multipara but it was not found significant. Parity was not found to associate with vitamin deficiency by other authors also.

Use of sunscreen lotion/cream was found to be associated with vitamin D deficiency. ANCs who has used sunscreen cream are more prone to developed vitamin D deficiency than who do not. This association with sunscreen cream also reported Holick MF et al¹⁵ in 1977 and Mallah et al¹⁶ 2011. Holick MF et al¹⁵ found that the most important source of vitamin D is skin's synthesis of vitamin D from sunlight and use of sun-block and time spend indoor increases the risk of vitamin D deficiency. Mallah et al¹⁶ 2011 conducted a study in Jordanians at national level revealed that women who wear hijab or niqab were more likely to have low levels vitamin D.

In this study, development of preeclampsia was not found to be associated with vitamin D deficiency. But Bodnar et al,¹⁷ Baker et al,¹⁸ Robinson et al¹⁹ and Rooplekha Chauhan¹² also found that prevalence of vitamin D deficiency is more prevalent in preeclamptic women.

In present study, LSCS were found significantly more in ANCs with vitamin D sufficiency whereas Dror D K et al²⁰ found no change in mean vitamin D level between women who underwent caesarean delivery and who did not. Moreover contrary to present findings, Merewood et al²¹ found that women with low vitamin D level had four times the odds of caesarean delivery.

In this study delivery of pre term babies was not found to be associated with vitamin D deficiency whereas Wagner et al²² in 2013 that supplemented 257 pregnant women since 12-16 week of gestational age with vitamin D (2000IU) found risk reduction in preterm infection and preterm birth.

In present study delivery of LBW babies was not found to be associated with vitamin D deficiency whereas Brook et al²³ also reported reduced incidence of low birth weight babies in vitamin D supplement Asian mother.

V. CONCLUSION

Vitamin D deficiency was found in 84.1% pregnant women. Religion and parity of women was not found to be associated with vitamin D deficiency whereas age, type of diet and use of sunscreen was found to be associated. Vitamin D deficiency was found significantly more in elderly, vegetarians and users of sunscreen than their counterparts.

Among maternal and neonatal outcomes, development of preeclampsia, LSCS as mode of delivery, delivery of pre term and LBW babies were having controversial findings. So it suggest much more researches in this regards.

CONFLICT OF INTEREST

None declared till now.

REFERENCES

- [1]. Holick MF, Chen TC. Vitamin D deficiency: A worldwide problem with health consequences. *Am J Clin Nutr* 2008;87:S1080-6.
- [2]. Van der wienen RP, Lowik MR, Van den berg H, De Groot LC, Haller J, Moreiras O et al. Serum vitamin D concentration among elderly people in Europe.1995;346:207-210.
- [3]. Pilz S, Dobnig H, Tomaschitz A, Kienreich K, Meinitzer A, Friedl C, dores W et al. Low 25-hydroxyvitamin D is associated with increased mortality in female nursing home residents. *J Clin Endocrinol Metab*. 2012 Apr;97(4):E653-657.
- [4]. Shibata M, Suzuki A, Sekiya T, Sekiguchi S, Asano S, Udagawa Y, et al. High prevalence of hypovitaminosis D in pregnant Japanese women with threatened premature delivery. *J Bone Miner Metab*. 2011 Sep;29(5):615–20.
- [5]. Harinarayan CV, Joshi SR. Vitamin D status in India--its implications and remedial measures. *J Assoc Physicians India*. 2009 Jan;57:40–8.
- [6]. Marwaha RK, Sripathy G. Vitamin D & bone mineral density of healthy school children in northern India. *Indian J Med Res*. 2008;127(3):239.
- [7]. Tandon VR, Sharma S, Mahajan S, Raina K, Mahajan A, Khajuria V, et al. Prevalence of vitamin d deficiency among Indian menopausal women and its correlation with diabetes: A first Indian cross sectional data. *J -Life Health*. 2014;5(3):121–5.
- [8]. Jani R, Palekar S, Munipally T, Ghugre P, Udipi S. Widespread 25-hydroxyvitamin D deficiency in affluent and nonaffluent pregnant Indian women. *BioMed Res Int*. 2014;2014:892162.
- [9]. Sachan A, Gupta R, Das V, Agarwal A, Awasthi PK, Bhatia V. High prevalence of vitamin D deficiency among pregnant women and their newborns in northern India. *Am J Clin Nutr*. 2005 May;81(5):1060–4.
- [10]. Dasgupta A, Saikia U, Sarma D. Status of 25(OH)D levels in pregnancy: A study from the North Eastern part of India. *Indian J Endocrinol Metab*. 2012 Dec;16(Suppl 2):S405-407
- [11]. Institute of medicine(IOM).Dietary reference intake for calcium,phosphorus,magnisium,vitamin D and fluoride.Washington (DC):National Academis;2011 dec 15
- [12]. Chauhan R, Chauhan M, Baghel P.prevelance of vitamin D deficiency and it's outcome in pregnancy at tertiary center.*International journal of medical and science*.2015;4 suppl 1(2320-3137):170-77
- [13]. Johnson DD, Wagner CL, Hulsey TC, McNeil RB, Ebeling M, Hollis BW. Vitamin D deficiency and insufficiency is common during pregnancy. *Am J Perinatol*. 2011 Jan;28(1):7–12.
- [14]. CroweFL,Steur M, Allen NE, Appleby PN, Travis RC, Key TJ. Plasma concentrations of 25-hydroxyvitamin D in meat eaters, fish eaters, vegetarians and vegans: results from the EPIC?Oxford study.*Public Health Nutr*. 2011 Feb;14(2):340–6
- [15]. Holick MF, Frommer JE, McNeill SC, Richtand NM, Henley JW, Potts JT. Photometabolism of 7-dehydrocholesterol to previtamin D3 in skin. *Biochem Biophys Res Commun*. 1977 May 9;76(1):107–14
- [16]. Mallah EM, Hamad MF, Elmanaseer MA, Qinna NA, Idkaidek NM, Arafat TA, et al. Plasma concentrations of 25-hydroxyvitamin D among Jordanians: Effect of biological and habitual factors on vitamin D status. *BMC Clin Pathol*. 2011 Aug 4;11:8
- [17]. Zhang C, Qiu C, Hu FB, David RM, van Dam RM, Bralley A, et al. Maternal plasma 25-hydroxyvitamin D concentrations and the risk for gestational diabetes mellitus. *PloS One*. 2008;3(11):e3753

-
- [18]. Baker AM, Haeri S, Camargo CA, Espinola JA, Stuebe AM. A nested case-control study of midgestation vitamin D deficiency and risk of severe preeclampsia. *J Clin Endocrinol Metab.* 2010 Nov;95(11):5105–9.
- [19]. Robinson CJ, Wagner CL, Hollis BW, Baatz JE, Johnson DD. Maternal vitamin D and fetal growth in early-onset severe preeclampsia. *Am J Obstet Gynecol.* 2011 Jun;204(6):556.e1-4
- [20]. Dror DK, King JC, Durand DJ, Allen LH. Association of modifiable and nonmodifiable factors with vitamin D status in pregnant women and neonates in Oakland, CA. *J Am Diet Assoc.* 2011 Jan;111(1):111–6
- [21]. Merewood A, Mehta SD, Chen TC, Bauchner H, Holick MF. Association between vitamin D deficiency and primary cesarean section. *J Clin Endocrinol Metab.* 2009 Mar;94(3):940–5
- [22]. Wagner CL, McNeil R, Hamilton SA, Winkler J, Cook CR, Warner G, et al. A randomized trial of vitamin D supplementation in 2 community health center networks in South Carolina. *Am J Obstet Gynecol.* 2013 Feb 1;208(2):137.e1-137.e13
- [23]. Brooke OG, Brown IR, Bone CD, Carter ND, Cleeve HJ, Maxwell JD, et al. Vitamin D supplements in pregnant Asian women: effects on calcium status and fetal growth. *Br Med J.* 1980 Mar 15;280(6216):751–4