

A Review of Smoking on Cardiovascular Disease: Impact on Human Health

Shashi K. Agarwal, MD

Center for Contemporary and Complementary Cardiology, 2227 US Highway 1, Suite 309, North Brunswick, NJ 08902, USA

Orcid # 0000-0003-0007-5582

Publons # E-4899-2016

Researchgate: https://www.researchgate.net/profile/Shashi_Agarwal

Received:- 8 March 2021/ Revised:- 16 March 2021/ Accepted: 21 March 2021/ Published: 31-03-2021

Copyright @ 2021 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— Cardiovascular diseases (CVD) are the leading cause of global morbidity and mortality. They also inflict huge direct and indirect financial costs on the worldwide society. With the ready availability of affordable therapeutics globally, and the relatively slow development of newer therapeutics, lifestyle behaviors are gaining importance, both as preventive and therapeutic interventions. Smoking tobacco is consistently and strongly related with a higher risk of CVD incidence, adverse progression, and increased mortality. Smoking is deleteriously associated with most cardiovascular diseases. Smokers die of more CVDs than from any other cause. Cessation of smoking is therefore paramount in improving worldwide CVD health. This paper reviews the negative effects of smoking on cardiovascular diseases.

Keywords— Smoking, Cardiovascular Diseases, Lifestyles, Modifiable Risk Factors, Nicotine.

I. INTRODUCTION

Tobacco is consumed by over a billion people around the world¹. Smoking is the most common mode of its use². This is done using products like cigarettes, e-cigarettes and water pipes³. Smoking tobacco exposes individuals to tobacco related noxious gaseous and particulate matter in several ways⁴⁻⁶. Most commonly, smoke enters the smoker's mouth directly from the cigarette being smoked and this is known as the first-hand smoke or mainstream smoke⁴. Side-stream cigarette smoke emanates from the burning ends of a cigarette and along with the exhaled main-stream smoke, results in second-hand smoke that could be inhaled by others⁵. Third-hand smoke is the residue from tobacco products that cling to surfaces such as skin, hair, clothing, and furniture⁶. These pollutants resist removal with traditional cleaning methods and may persist on these surfaces for several months. Tobacco is also consumed by non-smoking methods⁷. The smokeless tobacco products include loosely chewed tobacco leaves, tobacco paste and several tobacco-based concoctions such as naswar, gutka, and snuff⁸. The ingredients in these products are mainly absorbed via the mucous membranes of the mouth and nose.

Tobacco is highly addictive, primarily due to its nicotine content^{9,10}. All tobacco forms provide, besides nicotine, a plethora of harmful substances, many being extremely toxic and cancer provoking¹¹. These include many harmful nitrosamines and polycyclic aromatic hydrocarbons^{12,13}. Water pipe smoking results in large amounts of smoke being inhaled during a single smoking session and exposes the smoker to similar harmful substances as are present in cigarette smoke¹⁴. Inhaled e-cigarette vapor is usually lower in carcinogens and other toxins but is still harmful¹⁵. Smokeless tobacco also contains nicotine and more than 20 carcinogens, which enter the body via the oral and nasal mucous membranes¹⁶.

Tobacco smoking adversely affects literally every organ in the human body¹⁷. Tobacco smoke has a several thousand compounds, including several toxic ones, such as benzene, formaldehyde, benzopyrene, carbon monoxide, acrolein, and metals¹⁸. The respiratory tract is the first to be hit with these, (before disseminating all over the body via the blood stream) causing inflammation, irritation, asphyxiation, carcinogenesis, and other deleterious effects¹⁹. Further dissemination, mainly via the blood stream, results in the development of many other serious disorders including several forms of cancer²⁰. Smoking also affects pregnancy, causing a wide range of complications²¹. Infants exposed to cigarette smoke, either during the prenatal period or after birth have an increased risk of sudden death, orofacial clefts, periodontal disease, and dental caries²². Smoking accelerates aging, and smokers often develop wrinkles prematurely²³. Cigarette smoking greatly reduces

the quality of life²⁴. It is estimated that almost half of the smokers lose about 20 years of healthy life²⁵. In 2017, 182 million disability-adjusted life-years (DALYs) were attributed globally to tobacco use²⁶. Smokers over their lifetime demonstrate a higher mortality risk when compared to never smokers²⁷. According to the Global Burden of Disease Study, in 2015, worldwide, one out of ten deaths was due to smoking²⁸. Tobacco-related diseases are estimated to claim more lives than those due to AIDS, malaria and tuberculosis combined²⁹. The estimated economic cost associated with smoking is 1.8% of global gross domestic product³⁰.

Smoking cessation before the age of 40 is associated with a reduction of more than 90% of the excess morbidity and mortality³¹. Cessation at the age of 30 eliminates nearly all the excess risk. Besides bestowing a healthier life, cessation also increases life expectancy³¹. Jha et al estimated that smokers who started smoking early in adult life, gain almost 10, 9, 6, and 4 years of increased life if they stop smoking at ages of 30, 40, 50, or 60 years of age, respectively, compared to those who continue to smoke³².

II. DISCUSSION

The major CVDs are coronary heart disease (CHD) high blood pressure (HTN), stroke, heart failure (HF), cardiac arrhythmias, peripheral arterial disease (PAD), deep vein thrombosis (DVT) and vasculogenic erectile dysfunction (ED)³³. These impart the greatest non-communicable diseases burden globally and this is expected to rise globally in the coming years³⁴. They cause considerable disability and are responsible for an enormous number of deaths^{35,36}. They are responsible for a huge financial burden worldwide³⁴.

Smoking exerts adverse effects on several CVDs, resulting in poor outcomes³⁷. It is estimated that globally, smoking related CVD deaths exceed those from respiratory diseases or all forms of cancer combined³⁸. A smoker has approximately 10 years shorter lifespan than a nonsmoker³⁹. Smoking cessation is associated with a gradual reversion of the harmful cardiovascular effects⁴⁰. It reduces mortality and increases longevity⁴¹.

The relationship between smoking and CVD risk is non-linear⁴². It is estimated that 80% of the risk of smoking >20 cigarettes/day comes from smoking less than 3 cigarettes/day⁴³. Even smoking 1 cigarette/day is associated with 30–50% of the risk of CHD and 34–65% of the risk of stroke that would occur with smoking 20 cigarettes/day⁴⁴. Deleterious effects are also seen with low-tar cigarette, filtered cigarette, e-cigarette, cigar pipe, and water-pipe smoking⁴⁵. Nonsmokers exposed to passive smoke also demonstrate a 25–30% increased CV risk⁴⁶.

Tobacco smoking plays a critical role in the maintenance and development, and progression of atherosclerosis via several mechanisms⁴⁷. Tobacco smoking related chemicals increase low-density lipoprotein cholesterol levels, decrease high-density lipoprotein cholesterol level, increase inflammation, increase insulin resistance, increase catecholamines, increase the amount of fibrinogen, increase reactive oxygen species, and enhance platelet aggregation⁴⁷.

2.1 Smoking and HTN

According to the WHO, one in four men and one in five women have elevated BP globally and its prevalence is increasing steadily^{48,49}. Elevated BP increases the risk for coronary heart disease, PAD, stroke, kidney failure, and heart failure⁵⁰. It also increases CVD mortality⁵¹. It accounts for approximately 45% of all global CVD morbidity and mortality⁵².

The adverse relationship between smoking and hypertension is well documented⁵³. In the Physicians' Health Study, the risk of incident hypertension was increased by 15% in current smokers compared with never smokers⁵⁴. In a recent study, a more than 5 pack-years smoking exposure increased the risk of BP by more than 30%⁵⁵. Even passive smoking, after adjustments for several confounding factors, was found to be associated with a higher rate of hypertension in the exposed individuals, when compared to those not exposed to the secondhand cigarette smoke⁵⁶. Besides smoking promoted atherosclerosis, the sympathetic nervous system and the baroreflex-system also play a deleterious role⁵⁷⁻⁵⁹.

2.2 Smoking and CHD

CHD, or coronary artery disease/ischemic heart disease, is the most common CVD⁶⁰. Its prevalence is increasing in the middle- and low-income countries⁶¹. It is primarily caused by atherosclerosis, which leads to a flow-limiting stenosis of large epicardial coronary arteries, initially resulting in angina⁶². A plaque rupture or erosion can provoke superimposed atherothrombosis and vessel occlusion, leading to a myocardial infarction, or even death⁶³. It is responsible for an enormous amount of global disability⁴⁹. This imparts a heavy disability-adjusted life year (DALY) burden and smoking remains the number one CHD killer in the world^{49,65}.

Smoking is strongly linked with CAD⁶⁵. Smokers are two to four times more likely to develop CHD⁶⁶. In women, smokers between the ages of 35–39, almost triple their risk of CHD if they smoke just 1-4 cigarettes a day⁶⁷. The INTERHEART study found that smokers have a 1.95-fold increased risk of developing non-fatal myocardial infarction when compared to non-smokers⁶⁸. Hackshaw et al. reported that while smoking one cigarette per day increased the excess CHD risk by 40-50%, smoking 5-20 cigarettes per day increased the excess risk to only 55-65% - indicating that the smoking-cad relationship is non-linear⁶⁹. Passive smoking also increases the risk of ischemic heart disease⁷⁰. Smoking increases mortality⁷¹. One study showed that men smoking 20 cigarettes per day were twice as likely to die from CHD, when compared with never smokers⁷¹. Smoking cessation can reduce CHD mortality risk by around 36%⁷². This is impressive, when compared to a 29% reduction with statins, a 23% reduction with either beta-blockers or angiotensin-converting enzyme inhibitors, and a 15% reduction with aspirin⁷². Smoking cessation is therefore strongly recommended for all coronary artery disease patients by most major cardiology associations⁷³. The pathological basis of CHD is atherosclerosis, which is enhanced by smoking^{74,75}.

2.3 Smoking and Stroke

Stroke is characterized by a sudden onset of a neurological deficit due to a vascular cause and typically occurs after the age of 50 years⁷⁶. Vascular stroke may be ischemic or hemorrhagic or could result from a subarachnoid hemorrhage⁷⁶. Ischemic stroke is the most common⁷⁷. The Global Burden of Disease collaborators in a report published in 2018, estimated that the global lifetime risk of stroke from the age of 25 years was one in four and that the risk was twice as high for ischemic than hemorrhagic stroke⁷⁸. Stroke is associated with considerable functional disability⁷⁹. It is the second leading cause of death worldwide⁴⁹. The total number of strokes, the related DALY burden, and deaths, are steadily increasing globally, with the bulk occurring in the medium- and low-income world⁴⁹.

Smoking is a major risk factor for stroke⁸⁰, irrespective of the age of the individual^{81,82}. This increased risk appears to be dose-dependent⁸³, and the risk declines with smoking cessation⁸⁴. As noted with CHD, the risk is high even with smoking only one cigarette a day⁸⁵. Smoking one cigarette a day represents approximately 41% of the excess relative risk of getting a stroke when compared to the risk associated with smoking 20 cigarettes a day⁸⁵.

2.4 Smoking and Heart Failure

HF affects 2–4% of the global population, according to the World Health Organization⁸⁶. It is characterized by several functions restricting symptoms, such as shortness of breath, fatigue, and swelling of the ankles⁸⁷. These show frequent exacerbations and these result in repeated hospitalizations⁸⁷. The prognosis is extremely poor, and the mortality rate is high, with almost 50% of HF patients dying within 5 years⁸⁸. The main therapeutic goal in HF management is to reduce symptoms, decrease progression, diminish mortality, and improve the quality of life⁸⁹.

Smoking is associated with an increased risk of heart failure⁹⁰, and this is seen in both HFrEF (heart failure with reduced ejection fraction) and HFpEF (heart failure with preserved ejection fraction) after adjusting for other CVD risk factors⁹¹. The smoking-HF relationship is related to the amount and duration of smoking in HF patients⁹². Smokers not only demonstrate a reduced efficacy to treatment but are also more likely to experience undesirable long-term outcomes⁹³. Unfortunately, about 16% of HF patients continue to smoke⁹⁴. Exposure to second-hand smoke in HF patients is also detrimental and leads to a further reduction their quality of life⁹⁵. Active smoking increases mortality in HF patients⁹⁶. Even smoking in future donors results in a higher incidence of graft failure following heart transplantation and is associated with an increase in mortality⁹⁷. Smoking cessation is the leading modifiable risk factor for HF⁹⁸. The risk of HF declines steadily with increasing duration following smoking cessation⁹⁰.

The increased risk of HF and worse progression in existing HF in smokers are mediated via multifactorial pathways⁹⁹. These included increased inflammation, oxidative stress, impaired endothelial function, increased systemic vascular resistance, atherosclerosis, and ischemic heart disease¹⁰⁰⁻¹⁰³. They also demonstrate sympathetic predominance, increased pulmonary vascular resistance, toxic effects of inhaled carbon monoxide and harmful structural changes in the ventricles¹⁰⁴⁻¹⁰⁶.

2.5 Smoking and Arrhythmias

Atrial fibrillation (AF) is a common arrhythmic disorder¹⁰⁷. Its prevalence is higher in the elderly¹⁰⁸ and in individuals with other cardiovascular risk factors¹⁰⁹. It is a major risk factor for stroke, heart failure, and cognitive dysfunction, and the former two often being the first manifestations of AF¹¹⁰. Patients with established AF have a five-fold increase in their risk of stroke compared to those without this arrhythmia¹¹¹. It also reduces the individual's quality of life and is associated with increased mortality^{112,113}. According to Dai et al., AF contributed to about 290,000 deaths globally in 2017¹¹⁴.

The link between smoking and increased risk of developing AF is well established^{115,116}. A prospective, population-based study showed that the increased risk of developing atrial fibrillation was increased in both current and former cigarette smokers¹¹⁷. Similar conclusions have been reached in several other studies^{118,119}. Chamberlain et al found that current smoking increased the incidence of AF more than two-fold¹¹⁸. Former smokers also demonstrate a higher risk, although this risk is lower than that seen in current smokers¹¹⁹. Secondhand smoke exposure during gestational development and childhood also increases the risk of AF in later in life¹²⁰. Quitting smoking reduces the risk of AF¹²¹.

Even ventricular tachyarrhythmias are increased in smokers with ischemic left ventricular dysfunction¹²². Ventricular arrhythmias are a major cause of sudden cardiac death¹²³. Cigarette smoking is pro-arrhythmic due to the combined effects of nicotine, carbon monoxide, and pro-cyclic aromatic hydrocarbons^{124,125}.

2.6 Smoking and Valvular Diseases

Severe calcification of the aortic valve invariably results in impaired left ventricular outflow¹²⁶. The prevalence of moderate to severe aortic valve stenosis (AS) increases with age¹²⁶. Shortness of breath eventually develops in these patients and symptomology gradually progresses to angina, heart failure, and syncope¹²⁷. Once symptoms develop, mortality is extremely high, and more than half of AS patients die within 2 to 3 years¹²⁸. Treatment of severe aortic stenosis includes aortic valve replacement, either via a surgical or percutaneous approach¹²⁹. Globally, the prevalence of aortic valve calcification has steadily increased over the last 3 decades⁴⁹.

Several observational studies have established that smoking increases the risk of aortic valve stenosis¹³⁰⁻¹³³. Thanassoulis et al reported that greater cigarette use was associated with elevated odds for aortic valve calcification (AVC), even after adjusting for other cardiovascular factors¹³¹. In the MESA study, the risk for incident AVC was more than doubled in active smokers relative to never smokers, independent of other clinical factors¹³². The Cardiovascular Health Study reported 35% increased odds for aortic stenosis in smokers¹³³. This increased risk of aortic valve stenosis associated with smoking has also been noted in several prospective cohort studies¹³⁴. Stenotic aortic valve disease is pathologically like atherosclerosis, with lipoprotein deposition, chronic inflammation, and active leaflet calcification¹³⁵.

2.7 Smoking and Aortic Aneurysm

An aortic aneurysm, either thoracic or abdominal, is defined as a permanent localized dilatation of the vessel at least 150% compared to a relative normal adjacent diameter or with a maximal transverse diameter that exceeds 30 mm^{136,137}. It is estimated that abdominal aortic aneurysms, have a prevalence of 1.4% in those studied aged 50 to 84¹³⁸. They are potentially life-threatening, as rupture carries an extremely high mortality rate of 80–85%¹³⁹.

There is a strong association between smoking and the risk of developing abdominal aortic aneurysms, both in current and ex-smokers^{140,141}. It is estimated that smokers are almost 4 times more likely to die from aortic aneurysm than nonsmoking men and women¹⁴². Thoracic aortic aneurysms are uncommon but smoking also appears to increase the risk of their occurrence¹⁴³. Although decline in smoking has reduced their prevalence in some areas, abdominal aortic aneurysms continue to increase morbidity and mortality, in most parts of the world⁴⁹. Isolated aortic aneurysms develop from complex mechanisms, including apoptosis of vascular smooth muscle cells, oxidative stress, and inflammation¹⁴⁴.

2.8 Smoking and Peripheral Artery Disease

PAD is an atherosclerotic disease involving the pelvic and leg arteries¹⁴⁵. It is characterized by intermittent claudication¹⁴⁶. As the disease progresses, pain occurs at rest and the critical limb ischemia ultimately leads to tissue necrosis and amputation¹⁴⁷. PAD is a cardiovascular disease equivalent and is associated with a high rate of non-fatal and fatal cardiovascular events¹⁴⁸. PAD is diagnosed by measuring the ankle brachial index (ABI), and an ABI < 0.90 is extremely sensitive, and specific for a diagnosis of PAD^{149,150}. It affects more than 200 million people worldwide¹⁵¹. Its global prevalence and its associated mortality is rising⁴⁹.

Smoking is causally associated with PAD¹⁵². It is estimated that almost 80% of individuals with PAD smoke or have smoked in the past¹⁵³. Not only is the risk of PAD in smokers increased several-fold, but continuation of smoking also worsens prognosis in these individuals^{154,155}. Smoking is also associated with restenosis after endovascular revascularization, graft failure and amputation in PAD patients¹⁵⁶. Individuals smoking more than 1 pack per day have significantly greater adverse limb complications at one year, compared to those who smoked less¹⁵⁷. Continued smoking also increases the risk of myocardial infarctions and premature death in patients with PAD¹⁵⁸. Cessation of smoking helps reduce symptoms of

claudication in these patients but also helps reduce cardiovascular morbidity and mortality^{159,160}. PAD is an atherosclerotic disease and smoking promotes the pathogenesis of atherosclerosis, the primary pathology in these patients¹⁶¹.

2.9 Smoking and Erectile Dysfunction

ED is defined as a consistent or recurrent inability to attain or maintain a rigid penile erection suitable for satisfactory sexual intercourse¹⁶². It is a common male disorder and is expected to affect more than 320 million men worldwide by 2025^{163,164}. Causes may be endocrine or non-endocrine, and non-endocrine vasculogenic etiology is present in about 80% of cases¹⁶⁵. It is associated with many comorbidities and risk factors including increasing age, obesity (BMI>30 kg/m²), excessive alcohol intake, diabetes mellitus, other cardiovascular diseases, depression, and several other psychological disorders¹⁶⁶. It is usually a part of a more widespread atherosclerotic vascular disease and is known to be a strong predictor of other cardiovascular disorders, especially coronary artery disease¹⁶⁷. Its presence significantly reduces the quality of life in men and their partners¹⁶⁸.

Several epidemiological and prospective studies have shown that smoking increases ED^{169,170}. This adverse relationship has been noted in both former as well as current smokers¹⁷¹. He et al studied the association between smoking and ED among 7684 men without vascular disease¹⁷². The odds ratio of erectile dysfunction was 1.41 for cigarette smokers compared with never smokers¹⁷². Comparable data was obtained in men aged 40–70 years followed for up to 10 years, with the likelihood of developing ED being double in smokers¹⁷³. In a cross-sectional study of Chinese men in Hong Kong, smoking of ≥ 20 cigarettes daily was associated with a 50% increase of having ED¹⁷⁴. Smoking also increases the risk of ED in patients with other CVD risk factors¹⁷⁵⁻¹⁷⁷. The relationship between smoking and ED appears to be dose dependent¹⁷². In one study, penile rigidity during nocturnal erection inversely correlated with the number of cigarettes smoked per day¹⁷⁴. Smoking in ED is a modifiable risk factor and smoking cessation improves ED^{178,179}. This benefit has been noted in middle aged adults regardless of pack-year smoked¹⁸⁰.

Penile erection is largely caused by the presence of sufficient blood flow into the erectile tissue, simultaneous arterial endothelium-dependent dilatation, and sinusoidal endothelium-dependent corporal smooth muscle relaxation^{181,182}. This dilatation is impaired resulting in reduced blood flow to the penis in smokers, due to impaired nitric oxide (NO) synthesis and degradation¹⁸³. NO is the main neurotransmitter mediating vascular actions in the penile vasculature. Further, smoking results in the development of atherosclerosis in the internal pudendal and common penile arteries¹⁸⁴. Smoking may also cause ultrastructural damage to the corporal tissue¹⁸⁵.

2.10 Smoking and Venous Disease

DVT of the lower extremities is a common venous disorder and is associated with significant morbidity and a high rate of recurrence^{186,187}. These patients are at an extremely high risk of developing pulmonary embolism (PE)¹⁸⁸. PE is third most common cause of cardiovascular death in hospitalized patients in the West after stroke and heart attack¹⁸⁹. Acute PE is a significant cause of mortality worldwide, accounting for over 100,000 deaths in 2018 alone¹⁹⁰. Early diagnosis and intervention are paramount as 70% of deaths occur within the first hour of its occurrence¹⁹¹.

The increased risk for developing venous thrombo-embolism (VTE) with smoking has been determined by systemic meta-analytic studies^{192,193}. Kaptoge and group analyzed data from more than 1.1 million participants in 76 cohorts, and found that, smoking was associated with higher risk of VTE¹⁹⁴. El-Nasser recently re-confirmed this adverse link¹⁹⁵. Excess risk for VTE in smokers is also related to hospitalization¹⁹⁶. This risk is more during the first 12 weeks in smokers if there was surgery involved¹⁹⁷. Smoking induces endothelial dysfunction, along with oxidative stress and thrombo-inflammation and these contribute to the venous complications^{198,199}.

2.11 Smoking and other CVD risk factors

Smoking also aggravates many other CVD risk factors, such as diabetes mellitus²⁰⁰, dyslipidemia²⁰¹, insulin resistance²⁰², metabolic syndrome²⁰³, systemic inflammation²⁰⁴, and renal failure²⁰⁵.

III. CONCLUSION

The relationship between tobacco intake and the increased risk of CVDs is well established. The major mode of tobacco use is by smoking cigarettes. This is the number one preventable cause of CVDs. Further, quitting smoking after a major cardiovascular event is also the most effective secondary prevention measure²⁰⁶. The risks of CVDs drop dramatically following smoking cessation, irrespective of the age at cessation. Smoking cigarettes is also the number one preventable

cause of CVD mortality²⁰⁷. These risks are also increased with smokeless tobacco. Cessation of cigarette smoking dramatically reduces the risk of premature death from all smoking-related diseases²⁰⁸. The health risks associated with smoking show a dose-response relationship and there is no lower limit for deleterious effects²⁰⁹. No exposure to tobacco is safe for the cardiovascular system.

REFERENCES

- [1] <https://www.who.int/news-room/fact-sheets/detail/tobacco> - accessed February 13, 2021.
- [2] Smith PH, Assefa B, Kainth S, Salas-Ramirez KY, McKee SA, Giovino GA. Use of Mentholated Cigarettes and Likelihood of Smoking Cessation in the United States: A Meta-Analysis. *Nicotine Tob Res.* 2020;22(3):307-316. doi:10.1093/ntr/ntz067.
- [3] Shihadeh A, Schubert J, Klaiany J, El Sabban M, Luch A, Saliba NA. Toxicant content, physical properties and biological activity of waterpipe tobacco smoke and its tobacco-free alternatives. *BMJ.* 2015;24(1):22-30.
- [4] Pryor W.A., K Stone. Oxidants in cigarette smoke: Radicals, hydrogen peroxide, peroxyacetaldehyde, and peroxyacetylnitrate. *Ann NY Acad Sci,* 686 (1993), pp. 12-28.
- [5] A. E Taylor, D.C Johnson, H Kazemi. Environmental tobacco smoke and cardiovascular disease: A position paper from the Council on Cardiopulmonary and Critical Care, American Heart Association. *Circulation,* 86 (1992), pp. 699-702.
- [6] J.P. Winickoff, J. Friebely, S.E. Tanski, C. Sherrod, G.E. Matt, M.F. Hovell, et al. Beliefs about the health effects of “thirdhand” smoke and home smoking bans. *Pediatrics,* 123 (2009), pp. e74-e79.
- [7] Solhi M, Fattahi E, Barati H, Mohammadi M, Kasmaei P, Rastaghi S. Smokeless Tobacco Use in Iran: A Systematic Review. *Addict Health.* 2020;12(3):225-234. doi:10.22122/ahj.v12i3.274.
- [8] Kindvall M, Jonsson L, Schyberg M (2019) Freeze dried oral smokeless tobacco snuff or non-tobacco snuff product and method of manufacturing thereof. Retrieved from <https://patents.google.com/patent/US10278416B2/en>.
- [9] Sohn M, Hartley C, Froelicher ES, Benowitz NL. Tobacco use and dependence. *Semin Oncol Nurs.* 2003;19(4):250-260. doi:10.1053/j.soncn.2003.08.002.
- [10] J.T Powell. Vascular damage from smoking: Disease mechanisms at the arterial wall. *Vasc Med,* 3 (1998), pp. 21-28.
- [11] <https://www.cancer.org/cancer/cancer-causes/tobacco-and-cancer/carcinogens-found-in-tobacco-products.html>.
- [12] Konstantinou E, Fotopoulou F, Drosos A, et al. Tobacco-specific nitrosamines: A literature review. *Food Chem Toxicol.* 2018 Aug;118:198-203. doi: 10.1016/j.fct.2018.05.008.
- [13] Boström CE, Gerde P, Hanberg A, et al. Cancer risk assessment, indicators, and guidelines for polycyclic aromatic hydrocarbons in the ambient air. *Environ Health Perspect.* 2002 Jun;110 Suppl 3(Suppl 3):451-88. doi: 10.1289/ehp.110-1241197.
- [14] Eissenberg T, Shihadeh A. Waterpipe tobacco and cigarette smoking: direct comparison of toxicant exposure. *Am J Prev Med.* 2009;37:518-523.
- [15] R. Grana, N. Benowitz, S.A. Glantz. E-cigarettes: a scientific review. *Circulation,* 129 (2014), pp. 1972-1986.
- [16] Warnakulasuriya S, Straif K. Carcinogenicity of smokeless tobacco: evidence from studies in humans & experimental animals. *IJMR.* 2018;148(6):681.
- [17] https://www.cdc.gov/tobacco/data_statistics/sgr/2010/consumer_booklet/pdfs/consumer.pdf - accessed February 7, 2021.
- [18] <https://www.cancer.org/cancer/cancer-causes/tobacco-and-cancer/carcinogens-found-in-tobacco-products.html> – accessed February 2, 2021.
- [19] Lewis JB, Hirschi KM, Arroyo JA, Bikman BT, Kooyman DL, Reynolds PR. Plausible Roles for RAGE in Conditions Exacerbated by Direct and Indirect (Secondhand) Smoke Exposure. *International Journal of Molecular Sciences.* 2017; 18(3):652. <https://doi.org/10.3390/ijms18030652>
- [20] Lortet-Tieulent J., Sauer A.G., Siegel R.L., Miller K.D., Islami F., Fedewa S.A., Jacobs E.J., Jemal A. State-Level Cancer Mortality Attributable to Cigarette Smoking in the United States. *JAMA Intern. Med.* 2016;176:1792-1798. doi: 10.1001/jamainternmed.2016.6530.
- [21] O'Donnell MM, Baird J, Cooper C, et al. The Effects of Different Smoking Patterns in Pregnancy on Perinatal Outcomes in the Southampton Women's Survey. *Int J Environ Res Public Health.* 2020;17(21):7991. Published 2020 Oct 30. doi:10.3390/ijerph17217991
- [22] https://www.cdc.gov/reproductivehealth/maternalinfanthealth/substance-abuse/substance-abuse-during-pregnancy.htm?CDC_AA_refVal=https%3A%2F%2Fwww.cdc.gov%2Freproductivehealth%2Fmaternalinfanthealth%2Ftobaccousepregnancy%2Findex.htm – accessed February 12, 2021.
- [23] Clatici VG, Racoceanu D, Dalle C, Voicu C, Tomas-Aragones L, Marron SE, et al. Perceived age and lifestyle. The specific contributions of seven factors involved in health and beauty. *Maedica.* 2017;12(3):191-201.
- [24] Goldenberg M, Danovitch I, IsHak WW. Quality of life and smoking. *Am J Addict.* 2014 Nov-Dec;23(6):540-62. doi: 10.1111/j.1521-0391.2014.12148.x. Epub 2014 Sep 25. PMID: 25255868.
- [25] <http://www.kidon.com/smoke/percentages2.htm> - accessed January 22, 2021.
- [26] GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990-2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2018 Nov 10;392(10159):1923-1994. doi: 10.1016/S0140-6736(18)32225-6.

- [27] Inoue-Choi M, Liao LM, Reyes-Guzman C, Hartge P, Caporaso N, Freedman ND. Association of long-term, low-intensity smoking with all-cause and cause-specific mortality in the National Institutes of Health-AARP Diet and Health Study. *JAMA Intern Med.* 2017;177: 87–95. doi:10.1001/jamainternmed.2016.7511.
- [28] GBD 2015 Tobacco Collaborators MB, Fullman N, Ng M, et al. Smoking prevalence and attributable disease burden in 195 countries and territories, 1990–2015: a systematic analysis from the Global Burden of Disease Study 2015. *Lancet.* 2017;389:1885–1906. doi: 10.1016/S0140-6736(17)30819-X.
- [29] Fagerström K. The epidemiology of smoking: health consequences and benefits of cessation. *Drugs.* 2002;62(Suppl 2):1–9.
- [30] World No Tobacco Day 2017. Tobacco threatens us all: protect health, reduce poverty and promote development. Geneva: World Health Organization; 2017.
- [31] Jha P., Ramasundarahettige C., Landsman V., Rostron B., Thun M., Anderson R.N., McAfee T., Peto R. 21st-century hazards of smoking and benefits of cessation in the United States. *N. Engl. J. Med.* 2013;368:341–350. doi: 10.1056/NEJMsa1211128.
- [32] Jha P. The hazards of smoking and the benefits of cessation: a critical summation of the epidemiological evidence in high-income countries. *Elife.* 2020;9:e49979. Published 2020 Mar 24. doi:10.7554/eLife.49979.
- [33] https://www.who.int/cardiovascular_diseases/about_cvd/en/ - accessed October 29, 2020.
- [34] Bloom D, Cafiero E, Jané-Llopis E, Abrahams-Gessel S, Bloom L, et al. 2011. The Global Economic Burden of Non-Communicable Diseases. Geneva: World Econ. Forum.
- [35] Bourne RR. Global, regional, and national disability-adjusted life-years (DALYs) for 359 diseases and injuries and healthy life expectancy (HALE) for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet.* 2018;392:1859–1922.
- [36] Organization WHO. The top 10 causes of death. <https://www.who.int/news-room/fact-sheets/detail/the-top-10-causes-of-death>. Accessed 12 Apr 2020.
- [37] Mons U, Müezziner A, Gellert C, Schöttker B, Abnet CC, et al. CHANCES Consortium. Impact of smoking and smoking cessation on cardiovascular events and mortality among older adults: meta-analysis of individual participant data from prospective cohort studies of the CHANCES consortium. *BMJ.* 2015 Apr 20;350:h1551. doi: 10.1136/bmj.h1551.
- [38] Ezzati M, Lopez AD. Estimates of global mortality attributable to smoking in 2000. *Lancet* 362: 847–852, 2003. doi:10.1016/S0140-6736(03)14338-3.
- [39] Barua RS, Rigotti NA, Benowitz NL, Cummings KM, Jazayeri MA, Morris PB, et al. 2018 ACC expert consensus decision pathway on tobacco cessation treatment: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol.* 2018;72(25):3332–3365.
- [40] Ahmed AA, Patel K, Nyaku MA, et al. Risk of Heart Failure and Death After Prolonged Smoking Cessation: Role of Amount and Duration of Prior Smoking. *Circ Heart Fail.* 2015 Jul;8(4):694-701. doi: 10.1161/CIRCHEARTFAILURE.114.001885. Epub 2015 Jun 2. PMID: 26038535; PMCID: PMC5499230.
- [41] Mons Ute, MüezzinerAysel, Gellert Carolin, Schöttker Ben, Abnet Christian C, Bobak Martin et al. Impact of smoking and smoking cessation on cardiovascular events and mortality among older adults: meta-analysis of individual participant data from prospective cohort studies of the CHANCES consortium *BMJ* 2015; 350 :h1551
- [42] Doll R, Peto R, Boreham J, Sutherland I. Mortality in relation to smoking: 50 years' observations on male British doctors. *BMJ.* 2004;328(7455):1519.
- [43] Pope CA III, Burnett RT, Krewski D, Jerrett M, Shi Y, Calle EE, Thun MJ. Cardiovascular mortality and exposure to airborne fine particulate matter and cigarette smoke: shape of the exposure-response relationship. *Circulation* 120: 941–948, 2009. doi:10.1161/CIRCULATIONAHA.109.857888.
- [44] Hackshaw A, Morris JK, Boniface S, Tang JL, Milenković D. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports. *BMJ* 360: j5855, 2018. doi:10.1136/bmj.j5855.
- [45] He J, Vupputuri S, Allen K, Prerost MR, Hughes J, Whelton PK. Passive smoking and the risk of coronary heart disease—a meta-analysis of epidemiologic studies. *N Engl J Med.* 1999;340:920–926. doi: 10.1056/NEJM199903253401204.
- [46] Barua RS, Rigotti NA, Benowitz NL, Cummings KM, Jazayeri MA, Morris PB, et al. 2018 ACC expert consensus decision pathway on tobacco cessation treatment: a report of the American College of Cardiology Task Force on Clinical Expert Consensus Documents. *J Am Coll Cardiol.* 2018;72(25):3332–3365.
- [47] US Dept of Health and Human Services Introduction, evaluation of evidence on mechanisms of disease production, and summary. In: *How Tobacco Smoke Causes Disease: The Biology and Behavioral Basis for Smoking-Attributable Disease: A Report of the Surgeon General.* Published 2010. Accessed February 25, 2020. https://www.ncbi.nlm.nih.gov/books/NBK53017/pdf/Bookshelf_NBK53017.pdf.
- [48] WHO. A global brief on hypertension: Silent killer, global public health crisis (WHO/DCO/WHO/2013.2) World Health Organization; 2013. DOI: 10.5005/ijopmr-24-1-2.
- [49] Roth GA, Mensah GA, Johnson CO, et al.; GBD-NHLBI-JACC Global Burden of Cardiovascular Diseases Writing Group. Global Burden of Cardiovascular Diseases and Risk Factors, 1990-2019: Update From the GBD 2019 Study. *J Am Coll Cardiol.* 2020 Dec 22;76(25):2982-3021. doi: 10.1016/j.jacc.2020.11.010.
- [50] Whelton PK, Carey RM, Aronow WS, et al. ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA guideline for the prevention, detection, evaluation, and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Hypertension* 71 e13–e115, 2018. doi:10.1161/HYP.0000000000000065.

- [51] Institute for Health Metrics and Evaluation Global Health Data Exchange. 2016. [2019-09-10]. GBD Results Tool <http://ghdx.healthdata.org/gbd-results-tool>.
- [52] Costa SM, Lima CA, Nobre A, Vieira DMA, Leal, ALR. Hypertension bearers with high risk/big risk of cardiovascular diseases and socioeconomic and health indicators. *Rev Assoc Med Bras* (1992). 2018; 64(7): 601–10. DOI: 10.1590/1806-9282.64.07.601.
- [53] Dochi M, Sakata K, Oishi M, Tanaka K, Kobayashi E, Suwazono Y. Smoking as an independent risk factor for hypertension: a 14-year longitudinal study in male Japanese workers. *Tohoku J Exp Med*. 2009;217(1):37–43. doi: 10.1620/tjem.217.37.
- [54] Halperin RO, Gaziano JM, Sesso HD. Smoking and the risk of incident hypertension in middle-aged and older men. *Am J Hypertens* 21: 148–152, 2008. doi:10.1038/ajh.2007.36.
- [55] Kaplan RC, Baldoni PL, Strizich GM, et al. Current smoking raises risk of incident hypertension: Hispanic Community Health Study - Study of Latinos. *Am J Hypertens*. 2020 Sep 24:hpaa152. doi: 10.1093/ajh/hpaa152.
- [56] Li N, Li Z, Chen S, Yang N, Ren A, Ye R. Effects of passive smoking on hypertension in rural Chinese nonsmoking women. *J Hypertens*. 2015 Nov;33(11):2210-4. doi: 10.1097/HJH.0000000000000694. PMID: 26259123.
- [57] Yun M, Li S, Sun D, Ge S, Lai CC, Fernandez C, Chen W, Srinivasan SR, Berenson GS. Tobacco smoking strengthens the association of elevated blood pressure with arterial stiffness: the Bogalusa Heart Study. *J Hypertens* 33: 266–274, 2015. doi:10.1097/HJH.0000000000000410.
- [58] Cryer PE, Haymond MW, Santiago JV, Shah SD. Norepinephrine and epinephrine release and adrenergic mediation of smoking-associated hemodynamic and metabolic events. *N Engl J Med* 295: 573–577, 1976. doi:10.1056/NEJM197609092951101.
- [59] Pickering TG. Effects of stress and behavioral interventions in hypertension—the effects of smoking and nicotine replacement therapy on blood pressure. *J Clin Hypertens (Greenwich)* 3: 319–321, 2001. doi:10.1111/j.1524-6175.2001.00483.x.
- [60] Benjamin EJ, Muntner P, Alonso A, Bittencourt MS, Callaway CW, Carson AP, et al. Heart disease and stroke statistics—2019 update: a report from the American Heart Associationexternal icon. *Circulation*. 2019;139(10):e56–528.
- [61] Gaziano TA, Pagidipati N. Scaling up chronic disease prevention interventions in lower- and middle-income countries. *Annu Rev Public Health*. 2013;34:317–335. doi: 10.1146/annurev-publhealth-031912-114402.
- [62] Ganz P, Abben RP, Barry WH. Dynamic variations in resistance of coronary arterial narrowings in angina pectoris at rest. *Am J Cardiol*. 1987 Jan 1;59(1):66-70. doi: 10.1016/s0002-9149(87)80071-1.
- [63] Bentzon J.F., Otsuka F., Virmani R., Falk E. Mechanisms of plaque formation and rupture. *Circ. Res*. 2014;114:1852–1866. doi: 10.1161/CIRCRESAHA.114.302721.
- [64] Murray CJ, Lopez AD. Global mortality, disability, and the contribution of risk factors: Global Burden of Disease Study. *Lancet*. 1997 May 17;349(9063):1436-42. doi: 10.1016/S0140-6736(96)07495-8.
- [65] Le Bras A. Public health: light smoking and CVD risk. *Nat Rev Cardiol*. 2018;15(3):136. doi: 10.1038/nrcardio.2018.10.
- [66] Benjamin EJ, Virani SS, Callaway CW, et al. American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2018 Update: A Report From the American Heart Association. *Circulation* 137: e67–e492, 2018. doi:10.1161/CIR.0000000000000558.
- [67] Bjartveit K, Tverdal A. Health consequences of smoking 1-4 cigarettes per day. *Tob Control* 14: 315–320, 2005. doi:10.1136/tc.2005.011932.
- [68] Yusuf S, Hawken S, Ounpuu S, et al. Effect of potentially modifiable risk factors associated with myocardial infarction in 52 countries (the INTERHEART study): case-control study. *Lancet*. 2004;364:937–952. doi: 10.1016/S0140-6736(04)17018-9.
- [69] Hackshaw A, Morris JK, Boniface S, Tang JL, Milenković D. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports [published correction appears in *BMJ*. 2018 Apr 11;361:k1611. doi:10.1136/bmj.j5855.
- [70] Barnoya J, Glantz SA. Cardiovascular effects of secondhand smoke: nearly as large as smoking. *Circulation*. 2005;111:2684–2698. doi: 10.1161/circulationaha.104.492215.
- [71] Hammond EC, Horn D. Smoking and death rates--report on forty-four months of follow-up of 187,783 men. By E. Cuyler Hammond and Daniel horn, 1958. *CA Cancer J Clin*. 1988;38:28–58. doi: 10.3322/canjclin.38.1.28.
- [72] Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: a systematic review. *JAMA*. 2003; 290(1): 86–97. DOI: 10.1001/jama.290.1.86.
- [73] Ibanez B, James S, Agewall S, et al. 2017 ESC Guidelines for the management of acute myocardial infarction in patients presenting with ST-segment elevation: The Task Force for the management of acute myocardial infarction in patients presenting with ST-segment elevation of the European Society of Cardiology (ESC). *Eur Heart J*. 2018; 39(2): 119–77. DOI: 10.1093/eurheartj/ehx393.
- [74] Casolo G, Del Meglio J, Tessa C. Epidemiology and pathophysiologic insights of coronary atherosclerosis relevant for contemporary non-invasive imaging. *Cardiovasc Diagn Ther*. 2020 Dec;10(6):1906-1917. doi: 10.21037/cdt-20-157.
- [75] Salahuddin S, Prabhakaran D, Roy A. Pathophysiological mechanisms of tobacco-related CVD. *Glob Heart*. 2012;7:113–120. doi: 10.1016/j.gheart.2012.05.003.
- [76] Donnan GA, Fisher M, Macleod M, et al. Stroke. *Lancet* 2008;371:1612–23. 10.1016/S0140-6736(08)60694-7.
- [77] Barthels D, Das H. Current advances in ischemic stroke research and therapies. *BiochimBiophys Acta Mol Basis Dis*. 2020 Apr 1;1866(4):165260. doi: 10.1016/j.bbadis.2018.09.012.
- [78] Feigin VL, Nguyen G, Cercy K, Johnson CO, Alam T, Parmar PG, et al. . Global, regional, and country-specific lifetime risks of stroke, 1990 and 2016. *N Engl J Med*. (2018) 379:2429–37. 10.1056/NEJMoa1804492.

- [79] Verma R, Arya KN, Sharma P, Garg RK. Understanding gait control in post-stroke: implications for management. *J Bodyw Mov Ther.* 2012 Jan;16(1):14-21. doi: 10.1016/j.jbmt.2010.12.005.
- [80] Price AJ, Wright FL, Green J, Balkwill A, Kan SW, Yang TO, Floud S, Kroll ME, Simpson R, Sudlow CLM, Beral V, Reeves GK. Differences in risk factors for 3 types of stroke: UK prospective study and meta-analyses. *Neurology* 2018;90:e298–e306.
- [81] Mons U, Muezzinler A, Gellert C, Schottker B, Abnet CC, Bobak M, et al. Impact of smoking and smoking cessation on cardiovascular events and mortality among older adults: meta-analysis of individual participant data from prospective cohort studies of the CHANCES consortium. *BMJ.* 2015;350: h1551 10.1136/bmj.h1551.
- [82] Putaala J, Yesilot N, Waje-Andreassen U, Pitkaniemi J, Vassilopoulou S, Nardi K, et al. Demographic and geographic vascular risk factor differences in European young adults with ischemic stroke: the 15 cities young stroke study. *Stroke.* 2012;43: 2624–2630. 10.1161/STROKEAHA.112.662866.
- [83] Rissanen I, Oura P, Paananen M, Miettunen J, Geerlings MI. Smoking trajectories and risk of stroke until age of 50 years - The Northern Finland Birth Cohort 1966. *PLoS One.* 2019;14(12):e0225909. Published 2019 Dec 17. doi:10.1371/journal.pone.0225909.
- [84] Woodward M, Lam TH, Barzi F, Patel A, Gu D, Rodgers A, et al. Smoking, quitting, and the risk of cardiovascular disease among women and men in the Asia-Pacific region. *Int J Epidemiol.* 2005;34: 1036–1045. 10.1093/ije/dyi104.
- [85] Allan Hackshaw, Joan K Morris, Sadie Boniface, Jin-Ling Tang, Dušan Milenković. Low cigarette consumption and risk of coronary heart disease and stroke: meta-analysis of 141 cohort studies in 55 study reports. *BMJ* 2018; 360. doi: <https://doi.org/10.1136/bmj.j5855>.
- [86] WHO (World Health Organization Cardiovascular diseases (CVDs) [https://www.who.int/news-room/factsheets/detail/cardiovascular-diseases-\(cvds\)](https://www.who.int/news-room/factsheets/detail/cardiovascular-diseases-(cvds)). Published May 17, 2017. Accessed June 2, 2019.
- [87] Albano MG, Jourdain P, De Andrade V, Domenke A, Desnos M, d'Ivernois Jean-François. Therapeutic patient education in heart failure: do studies provide sufficient information about the educational programme? *Arch Cardiovasc Dis.* 2014;107(5):328–339. doi: 10.1016/j.acvd.2013.12.002.
- [88] Benjamin EJ, Blaha MJ, Chiuve SE, Cushman M, Das SR, Deo R, de Ferranti SD, Floyd J, Fornage M, Gillespie C, Isasi CR, et al. American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2017 Update: A Report From the American Heart Association. *Circulation.* 2017 Mar 7;135(10):e146-e603. doi: 10.1161/CIR.0000000000000485.
- [89] Inamdar AA, Inamdar AC. Heart Failure: Diagnosis, Management and Utilization. *J Clin Med.* 2016;5(7) doi: 10.3390/jcm5070062.
- [90] Aune D, Schlesinger S, Norat T, Riboli E. Tobacco smoking and the risk of heart failure: a systematic review and meta-analysis of prospective studies. *Eur J PrevCardiol* 2019;26:279–288.
- [91] Chambless LE, Heiss G, Folsom AR, Rosamond W, Szklo M, Sharrett AR, Clegg LX. Association of coronary heart disease incidence with carotid arterial wall thickness and major risk factors: the Atherosclerosis Risk in Communities (ARIC) Study, 1987–1993. *American journal of epidemiology* 1997;146:483–494.
- [92] Feodoroff M, Harjutsalo V, Forsblom C, Groop PH. Dose-dependent effect of smoking on risk of coronary heart disease, heart failure and stroke in individuals with type 1 diabetes. *Diabetologia.* 2018;61(12):2580–2589. doi: 10.1007/s00125-018-4725-9.
- [93] Ambrose JA, Barus RS. The pathophysiology of cigarette smoking and cardiovascular disease. *J Am Coll Cardiol.* 2004;43(10):1731–1737. doi: 10.1016/j.jacc.2003.12.047.
- [94] Conard MW, Haddock CK, Poston WS, Spertus JA. The impact of smoking status on the health status of heart failure patients. *Congest Heart Fail.* 2009;15(2):82–86. doi: 10.1111/j.1751-7133.2009.00053.x.
- [95] Weeks SG, Glantz SA, De Marco T, Rosen AB, Fleischmann KE. Secondhand smoke exposure and quality of life in patients with heart failure. *Arch Intern Med.* 2011;171:1887–1893. doi: 10.1001/archinternmed.2011.518.
- [96] Suskin N, Sheth T, Negassa A, Yusuf S. Relationship of current and past smoking to mortality and morbidity in patients with left ventricular dysfunction. *J Am Coll Cardiol.* 2001;37:1677–1682. doi: 10.1016/S0735-1097(01)01195-0.
- [97] Hussain Z, Yu M, Wozniak A, Kim D, Krepostman N, Liebo M, Raichlin E, Heroux A, Joyce C, Ilias-Basha H. Impact of Donor Smoking History on Post Heart Transplant Outcomes: A Propensity Matched Analysis of ISHLT registry. *Clin Transplant.* 2020 Oct 23:e14127. doi: 10.1111/ctr.14127.
- [98] Fleg JL. Healthy lifestyle and risk of heart failure: An ounce of prevention well worth the effort. *Circ Heart Fail.* 2016;9(4):e003155. doi: 10.1161/CIRCHEARTFAILURE.116.003155.
- [99] Centers for Disease C, Prevention, National Center for Chronic Disease P, et al. Publications and reports of the Surgeon General. How tobacco smoke causes disease: the biology and behavioral basis for smoking-attributable disease: a report of the surgeon general, Centers for Disease Control and Prevention (US), Atlanta (GA) (2010).
- [100] Anker SD, von Haehling S. Inflammatory mediators in chronic heart failure: an overview. *Heart.* 2004;90:464–470. doi: 10.1136/hrt.2002.007005.
- [101] Munzel T, Gori T, Keaney JJ, Maack C, Daiber A. Pathophysiological role of oxidative stress in systolic and diastolic heart failure and its therapeutic implications. *Eur Heart J.* 2015;36:2555–2564. doi: 10.1093/eurheartj/ehv305.
- [102] Morris PB, Ference BA, Jahangir E, et al. Cardiovascular effects of exposure to cigarette smoke and electronic cigarettes: clinical perspectives from the Prevention of Cardiovascular Disease Section Leadership Council and Early Career Councils of the American College of Cardiology. *J Am Coll Cardiol.* 2015; 66:1378–1391. doi: 10.1016/j.jacc.2015.07.037.
- [103] Kamimura D, Cain LR, Mentz RJ, et al. Cigarette smoking and incident heart failure. *Circulation.* 2018;137(24):2572–2582. doi: 10.1161/CIRCULATIONAHA.117.031912.

- [104] Nicolozakes AW, Binkley PF, Leier CV. Hemodynamic effects of smoking in congestive heart failure. *Am J Med. Sci.* 1988; 296:377–380.
- [105] Reboul C, Boissière J, André L, Meyer G, Bideaux P, Fouret G, Feillet-Coudray C, Obert P, Lacampagne A, Thireau J, Cazorla O, Richard S. Carbon monoxide pollution aggravates ischemic heart failure through oxidative stress pathway. *Sci Rep.* 2017; 7:39715. doi: 10.1038/srep39715.
- [106] Ambrose JA, Barus RS. The pathophysiology of cigarette smoking and cardiovascular disease. *J Am Coll Cardiol.* 2004;43(10):1731–1737. doi: 10.1016/j.jacc.2003.12.047.
- [107] Krijthe BP, Kunst A, Benjamin EJ, Lip GY, Franco OH, Hofman A, et al. Projections on the number of individuals with atrial fibrillation in the European Union, from 2000 to 2060. *Eur Heart J.* 2013;34(35):2746–51.
- [108] Go AS, Hylek EM, Phillips KA, Chang Y, Henault LE, Selby JV, et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. *JAMA.* 2001;285(18):2370–5. doi: http://dx.doi.org/10.1001/jama.285.18.2370.
- [109] Staerk L, Wang B, Preis SR, Larson MG, Lubitz SA, Ellinor PT, et al. Lifetime risk of atrial fibrillation according to optimal, borderline, or elevated levels of risk factors: cohort study based on longitudinal data from the Framingham Heart Study. *BMJ.* 2018;361:k1453. doi: http://dx.doi.org/10.1136/bmj.k1453.
- [110] Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke.* 1991;22(8):983–8. doi: http://dx.doi.org/10.1161/01.STR.22.8.983.
- [111] Conen D, Chae CU, Glynn RJ, Tedrow UB, Everett BM, Buring JE, et al. Risk of death and cardiovascular events in initially healthy women with new-onset atrial fibrillation. *JAMA.* 2011;305(20):2080–7. doi: http://dx.doi.org/10.1001/jama.2011.659.
- [112] Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke.* 1991;22(8):983–8.
- [113] Conen D, Chae CU, Glynn RJ, Tedrow UB, Everett BM, Buring JE, et al. Risk of death and cardiovascular events in initially healthy women with new-onset atrial fibrillation. *JAMA.* 2011;305(20):2080–7. doi: http://dx.doi.org/10.1001/jama.2011.659.
- [114] Dai H, Zhang Q, Much AA, Maor E, Segev A, Beinart R, Adawi S, Lu Y, Bragazzi NL, Wu J. Global, regional, and national prevalence, incidence, mortality, and risk factors for atrial fibrillation, 1990–2017: results from the Global Burden of Disease Study 2017. *Eur Heart J Qual Care Clin Outcomes* 2020;doi: 10.1093/ehjqcco/qcaa061.
- [115] Rahman F, Yin X, Larson MG, Ellinor PT, Lubitz SA, Vasani RS, McManus DD, Magnani JW, Benjamin EJ. Trajectories of risk factors and risk of new-onset atrial fibrillation in the Framingham Heart Study. *Hypertension* 68: 597–605, 2016. doi:10.1161/HYPERTENSIONAHA.116.07683.
- [116] Zuo H, Nygård O, Vollset SE, Ueland PM, Ulvik A, Midttun Ø, Meyer K, Igland J, Sulo G, Tell GS. Smoking, plasma cotinine and risk of atrial fibrillation: the Hordaland Health Study. *J Intern Med.* 2018 Jan;283(1):73–82. doi: 10.1111/joim.12689.
- [117] Heeringa J, Kors JA, Hofman A, van Rooij FJ, Witteman JC. Cigarette smoking and risk of atrial fibrillation: the Rotterdam Study. *Am Heart J.* 2008 Dec;156(6):1163–9. doi: 10.1016/j.ahj.2008.08.003.
- [118] Chamberlain AM, Agarwal SK, Folsom AR, Duval S, Soliman EZ, Ambrose M, Eberly LE, Alonso A. Smoking and incidence of atrial fibrillation: results from the Atherosclerosis Risk in Communities (ARIC) study. *Heart Rhythm.* 2011 Aug;8(8):1160–6. doi: 10.1016/j.hrthm.2011.03.038.
- [119] Aune D, Schlesinger S, Norat T, Riboli E. Tobacco smoking and the risk of atrial fibrillation: A systematic review and meta-analysis of prospective studies. *Eur J Prev Cardiol.* 2018 Sep;25(13):1437–1451. doi: 10.1177/2047487318780435.
- [120] S. Dixit, M.J. Pletcher, E. Vittinghoff, et al. Secondhand smoke and atrial fibrillation: data from the Health eHeart Study. *Heart Rhythm*, 13 (2016), pp. 3–9.
- [121] S. Suzuki, T. Otsuka, K. Sagara, et al. Association between smoking habits and the first-time appearance of atrial fibrillation in Japanese patients: evidence from the Shinken Database. *J Cardiol*, 66 (2015), pp. 73–79.
- [122] Goldenberg I, Moss AJ, McNitt S, Zareba W, Daubert JP, Hall WJ, et al. Cigarette smoking and the risk of supraventricular and ventricular tachyarrhythmias in high-risk cardiac patients with implantable cardioverter defibrillators. *J Cardiovasc Electrophysiol.* 2006;17(9):931–936.
- [123] Messerli FH, Nunez BD, Ventura HO, Snyder DW. Overweight and sudden death: increased ventricular ectopy in cardiomyopathy of obesity. *Arch Intern Med*, 147 (1987), pp. 1725–1728.
- [124] Benowitz NL. Drug therapy. Pharmacologic aspects of cigarette smoking and nicotine addiction. *N Engl J Med.* 1988;319(20):1318–1330.
- [125] Howard G, Wagenknecht LE, Burke GL, Diez-Roux A, Evans GW, McGovern P, et al. Cigarette smoking and progression of atherosclerosis: the Atherosclerosis Risk in Communities (ARIC) study. *JAMA.* 1998;279(2):119–124.
- [126] Osnabrugge RL, Mylotte D, Head SJ, Van Mieghem NM, Nkomo VT, LeReun CM, et al. Aortic stenosis in the elderly: disease prevalence and number of candidates for transcatheter aortic valve replacement: a meta-analysis and modeling study. *J Am Coll Cardiol.* 2013;62(11):1002–12.
- [127] Bottaro G, Zappulla P, Deste W, Famà F, Agnello F, Trovato D, Indelicato A, Barbanti M, Sgroi C, Monte IP, Tamburino C. Severe Aortic Valve Stenosis: Symptoms, Biochemical Markers, and Global Longitudinal Strain. *J Cardiovasc Echogr.* 2020 Jul-Sep;30(3):154–161. doi: 10.4103/jcecho.jcecho_13_20.
- [128] Liu Z, Duarte R, Kidney E, Bem D, Bramley G, Rooney S, et al. Clinical effectiveness and safety of transcatheter aortic valve implantation (TAVI) for aortic stenosis: a systematic review and meta-analysis [Internet]. Birmingham (UK): University of Birmingham; 2016.

- [129] Takeji Y, Taniguchi T, Morimoto T, et al. the CURRENT AS Registry Investigators and K-TAVI Registry Investigators. Transcatheter Aortic Valve Implantation vs. Surgical Aortic Valve Replacement for Severe Aortic Stenosis in Real-World Clinical Practice. *Circ J*. 2020 Apr 24;84(5):806-814. doi: 10.1253/circj.CJ-19-0951.
- [130] Larsson SC, Wolk A, Back M.. Alcohol consumption, cigarette smoking and incidence of aortic valve stenosis. *J Intern Med* 2017;282:332–339.
- [131] Thanassoulis G, Massaro JM, Cury R et al. Associations of long-term and early adult atherosclerosis risk factors with aortic and mitral valve calcium. *J Am Coll Cardiol* 2010;55:2491–8.
- [132] Owens DS, Katz R, Takasu J, Kronmal R, Budoff MJ, O'Brien KD. Incidence and progression of aortic valve calcium in the Multi-ethnic Study of Atherosclerosis (MESA). *Am J Cardiol* 2010;105:701–8.
- [133] Stewart BF, Siscovick D, Lind BK et al. Clinical factors associated with calcific aortic valve disease. *J Am Coll Cardiol* 1997;29:630–4.
- [134] Martinsson A, Ostling G, Persson M, Sundquist K, Andersson C, Melander O, Engstrom G, Hedblad B, Smith JG.. Carotid plaque, intima-media thickness, and incident aortic stenosis: a prospective cohort study. *ArteriosclerThrombVasc Biol* 2014;34:2343–2348.
- [135] Lester SJ, Heilbron B, Gin K, Dodek A, Jue J. The natural history and rate of progression of aortic stenosis. *Chest*. 1998;113(4):1109–14. 10.1378/chest.113.4.1109.
- [136] Johnston KW, Rutherford RB, Tilson MD, Shah DM, Hollier L, Stanley JC. Suggested standards for reporting on arterial aneurysms. Subcommittee on Reporting Standards for Arterial Aneurysms, Ad Hoc Committee on Reporting Standards, Society for Vascular Surgery and North American Chapter, International Society for Cardiovascular Surgery. *J Vasc Surg*. 1991 Mar;13(3):452-8.
- [137] Lim J., Wolff J., Rodd C., Cooper D., Earnshaw J.J. Outcome in Men with a Screen-detected Abdominal Aortic Aneurysm Who are not Fit for Intervention. *Eur. J. Vasc. Endovasc. Surg*. 2015;50:732–736. doi: 10.1016/j.ejvs.2015.07.035.
- [138] Kent KC, Zwolak RM, Egorova NN, Riles TS, Manganaro A, Moskowitz AJ, Gelijns AC, Greco G. Analysis of risk factors for abdominal aortic aneurysm in a cohort of more than 3 million individuals. *J Vasc Surg*. 2010 Sep;52(3):539-48.
- [139] Budd J.S., Finch D., Carter P. A study of the mortality from ruptured abdominal aortic aneurysms in a district community. *Eur. J. Vasc. Surg*. 1989;3:351–354. doi: 10.1016/S0950-821X(89)80073-8.
- [140] Aune D, Schlesinger S, Norat T, Riboli E.. Tobacco smoking and the risk of abdominal aortic aneurysm: a systematic review and meta-analysis of prospective studies. *Sci Rep* 2018;8:14786.
- [141] NICE Guideline Updates Team (UK). Risk factors for predicting presence of an abdominal aortic aneurysm: Abdominal aortic aneurysm: diagnosis and management: Evidence review A. London: National Institute for Health and Care Excellence (UK); 2020 Mar. PMID: 32407017.
- [142] United States Department of Health and Human Services. The Health Consequences of Smoking: 50 Years of Progress. A Report of the Surgeon General. Atlanta, GA: Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2014.
- [143] Landenhed M, Engstrom G, Gottsater A, Caulfield MP, Hedblad B, Newton-Cheh C, Melander O, Smith JG.. Risk profiles for aortic dissection and ruptured or surgically treated aneurysms: a prospective cohort study. *J Am Heart Assoc* 2015;4:e001513.
- [144] Kopacz A, Werner E, Grochot-Przędzek A, et al. Simvastatin Attenuates Abdominal Aortic Aneurysm Formation Favoured by Lack of Nrf2 Transcriptional Activity. *Oxid Med Cell Longev*. 2020;2020:6340190. Published 2020 Jun 16. doi:10.1155/2020/6340190.
- [145] Fowkes FG, Murray GD, Butcher I, et al. Ankle brachial index combined with Framingham Risk Score to predict cardiovascular events and mortality: a meta-analysis. *JAMA*. 2008;300(2):197–208.
- [146] Marashi M, Bahr A, Al Mahdi H, Bakir S, Al Farhan M. Global Risk Profile Verification in Patients with Venous Thromboembolism (GRIP VTE) in 5 Gulf countries. *Clin Appl ThrombHemost*. 2009 May-Jun;15(3):289-96. doi: 10.1177/1076029608315168.
- [147] Caitlin W. Hicks, Chao Yang, Chiadi E. et al. Association s of Obesity With Incident Hospitalization Related to Peripheral Artery Disease and Critical Limb Ischemia in the ARIC Study. *Journal of the American Heart Association*. 2018;7. 9 Aug 2018https://doi.org/10.1161/JAHA.118.008644.
- [148] Subherwal S, Patel MR, Kober L, Peterson ED, Bhatt DL, Gislason GH, Olsen AM, Jones WS, Torp-Pedersen C, Fosbol EL. Peripheral artery disease is a coronary heart disease risk equivalent among both men and women: results from a nationwide study. *Eur J PrevCardiol*. 2015 Mar;22(3):317-25.
- [149] Gerhard-Herman MD, Gornik HL, Barrett C, et al. 2016 AHA/ACC Guideline on the Management of Patients With Lower Extremity Peripheral Artery Disease: Executive Summary: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *J Am Coll Cardiol*. 2017 Mar 21;69(11):1465-1508. doi: 10.1016/j.jacc.2016.11.008.
- [150] Aboyans V, Ricco JB, Bartelink MEL, et al. Desormais I. 2017 ESC Guidelines on the Diagnosis and Treatment of Peripheral Arterial Diseases, in collaboration with the European Society for Vascular Surgery (ESVS): document covering atherosclerotic disease of extracranial carotid and vertebral, mesenteric, renal, upper and lower extremity arteries. Endorsed by: the European Stroke Organization (ESO)The Task Force for the Diagnosis and Treatment of Peripheral Arterial Diseases of the European Society of Cardiology (ESC) and of the European Society for Vascular Surgery (ESVS) *Eur Heart J*. 2018;39:763–816. doi: 10.1093/eurheartj/ehx095.
- [151] Alzamora MT, Fore's R, Baena-Dí'ez JM, Pera G, Toran P, Sorribes M et al (2010) The peripheral arterial disease study (PERART/ARTPER): prevalence and risk factors in the general population. *BMC Public Health* 10:38.
- [152] Lu L, Mackay DF, Pell JP. Meta-analysis of the association between cigarette smoking and peripheral arterial disease. *Heart* 2014;100:414–423.
- [153] Smith GD, Shipley MJ, Rose G. Intermittent claudication, heart disease risk factors, and mortality. The Whitehall Study. *Circulation* 1990;82:1925-31.

- [154] Benjamin EJ, Virani SS, Callaway CW, et al. American Heart Association Council on Epidemiology and Prevention Statistics Committee and Stroke Statistics Subcommittee. Heart Disease and Stroke Statistics-2018 Update: A Report From the American Heart Association. *Circulation* 137: e67–e492, 2018. doi:10.1161/CIR.0000000000000558.
- [155] Aboyans V, Criqui MH, Denenberg JO, Knokke JD, Ridker PM, Fronck A. Risk factors for progression of peripheral arterial disease in large and small vessels. *Circulation*. 2006 Jun 6;113(22):2623-9. doi: 10.1161/CIRCULATIONAHA.105.608679.
- [156] Willigendael EM, Teijink JAW, Bartelink ML, Peters RJG, Büller HR, Prins MH. Smoking and the patency of lower extremity bypass grafts: a meta-analysis. *J Vasc Surg* 2005;42:67-74.
- [157] Young JC, Paul NJ, Karatas TB, Kondrasov SA, McGinagle KL, Crouner JR, Pascarella L, Farber MA, Kibbe MR, Marston WA, Kalbaugh CA. Cigarette smoking intensity informs outcomes after open revascularization for peripheral artery disease. *J Vasc Surg*. 2019 Dec;70(6):1973-1983.e5. doi: 10.1016/j.jvs.2019.02.066.
- [158] Price JF, Mowbray PI, Lee AJ, Rumley A, Lowe GD, Fowkes FG. Relationship between smoking and cardiovascular risk factors in the development of peripheral arterial disease and coronary artery disease: Edinburgh Artery Study. *Eur Heart J*. 1999 Mar;20(5):344-53. doi: 10.1053/euhj.1998.1194.
- [159] Ratchford EV. Medical management of claudication. *J Vasc Surg* 2017;66:275-80.
- [160] Lu JT, Creager MA. The relationship of cigarette smoking to peripheral arterial disease. *Rev Cardiovasc Med*. 2004 Fall;5(4):189-93. PMID: 15580157.
- [161] Centner AM, Bhide PG, Salazar G. Nicotine in Senescence and Atherosclerosis. *Cells*. 2020 Apr 22;9(4):1035. doi: 10.3390/cells9041035.
- [162] Muneer A, Kalsi J, Nazareth I, Arya M. Erectile dysfunction. *BMJ*. 2014 Jan 27;348:g129.
- [163] NIH Consensus Conference. Impotence. NIH Consensus Development Panel on Impotence. *JAMA* 1993;270:83-90. Doi: 10.1001/jama.1993.03510010089036.
- [164] Aytia IA, McKinlay JB, Krane RJ. The likely worldwide increase in erectile dysfunction between 1995 and 2025 and some possible policy consequences. *BJU Int* 1999;84:50-6. 10.1046/j.1464-410x.1999.00142.x.
- [165] Yafi FA, Jenkins L, Albersen M, Corona G, Isidori AM, Goldfarb S, et al. Erectile dysfunction. *Nat Rev Dis Primers*. 2016;2:16003.
- [166] Kim S, Cho MC, Cho SY, Chung H, Rajasekaran MR. Novel Emerging Therapies for Erectile Dysfunction. *World J Mens Health*. 2021 Jan;39(1):48-64. doi: 10.5534/wjmh.200007.
- [167] Bryan G, Schwartz, Robert A. Kloner. Cardiovascular Implications of Erectile Dysfunction. *Circulation*. 2011;123:e609–e611.
- [168] Mulhall JP, Giraldo A, Hackett G, et al. The 2018 Revision to the Process of Care Model for Evaluation of Erectile Dysfunction. *J Sex Med* 2018;15:1280-92. 10.1016/j.jsxm.2018.06.005.
- [169] Millett C, Wen LM, Rissel C, Smith A, Richters J, Grulich A, de Visser R. Smoking and erectile dysfunction: Findings from a representative sample of Australian men. *Tob Control*. 2006;15:136–139.
- [170] Bacon CG, et al. A prospective study of risk factors for erectile dysfunction. *J Urol*. 2006;176:217–221.
- [171] Gades NM, Nehra A, Jacobson DJ, McGree ME, Girman CJ, Rhodes T, Roberts RO, Lieber MM, Jacobsen SJ. Association between smoking and erectile dysfunction: A population-based study. *Am J Epidemiol*. 2005;161:346–351.
- [172] He J, Reynolds K, Chen J, et al. Cigarette smoking and erectile dysfunction among Chinese men without clinical vascular disease. *Am J Epidemiol*. 2007;166(7):803–809. doi:10.1093/aje/kwm154.
- [173] Feldman HA, Johannes CB, Derby CA, Kleinman KP, Mohr BA, Araujo AB, McKinlay JB. Erectile dysfunction and coronary risk factors: Prospective results from the Massachusetts Male Aging Study. *Prev Med*. 2000;30:328–338.
- [174] Lam TH, Abdullah AS, Ho LM, et al. Smoking and sexual dysfunction in Chinese males: findings from men's health survey. *Int J Impot Res* 2006;18:364-9. 10.1038/sj.ijir.3901436.
- [175] Shabsigh R, Fishman IJ, Schum C, Dunn JK. Cigarette smoking and other vascular risk factors in vasculogenic impotence. *Urology*. 1991;38:227–231.
- [176] Bortolotti A, Fedele D, Chatenoud L, Colli E, Coscelli C, Landoni M, Lavezzari M, Santeusano F, Parazzini F. Cigarette smoking: A risk factor for erectile dysfunction in diabetics. *Eur Urol*. 2001;40:392–396.
- [177] Azab SS, Salem A, Ismail NN, El Khat Y, El Gebally MA. Penile hemodynamics study in erectile dysfunction men: the influence of smoking obesity on the parameters of penile duplex. *Int Urol Nephrol*. 2020 Jun;52(6):1015-1025. doi: 10.1007/s11255-020-02405-1.
- [178] Shamloul R, Ghanem H. Erectile dysfunction. *Lancet* 2013;381:153-65. 10.1016/S0140-6736(12)60520-0.
- [179] Pourmand G, Alidaee MR, Rasuli S, et al. Do cigarette smokers with erectile dysfunction benefit from stopping?: a prospective study. *BJU Int* 2004;94:1310-3. 10.1111/j.1464-410X.2004.05162.x.
- [180] Sahin MO, Sen V, Gunduz G, Ucer O. Effect of smoking cessation on sexual functions in men aged 30 to 60 years. *Int Braz J Urol*. 2020 Jul-Aug;46(4):642-648. doi: 10.1590/S1677-5538.IBJU.2019.0541.
- [181] Lewis RW, Hatzichristou DG, Laumann E, McKinlay JB. Epidemiology and natural history of erectile dysfunction; risk factors including iatrogenic and aging. In: Jardin A, Wagner AH, Khoury S, Guiliano F, Padma-Nathan H, Rosen A, eds, editors. *Erectile Dysfunction*. Plymouth: Health Publication Ltd; 2000. pp. 21–51.
- [182] Powell JT. Vascular damage from smoking: disease mechanisms at the arterial wall. *Vasc Med*. 1998;3:21–28.6.
- [183] Tostes RC, Carneiro FS, Lee AJ, et al. Cigarette smoking and erectile dysfunction: focus on NO bioavailability and ROS generation. *J Sex Med*. 2008;5(6):1284-1295. doi:10.1111/j.1743-6109.2008.00804.x].

- [184] Rosen MP, Greenfield AJ, Walker TG, Grant P, Dubrow J, Bettmann MA, et al. Cigarette smoking: an independent risk factor for atherosclerosis in the hypogastric-cavernous arterial bed of men with arteriogenic impotence. *J Urol*. 1991;145:759-63.7,
- [185] Rehill N, Beck CR, Yeo KR, Yeo WW. The effect of chronic tobacco smoking on arterial stiffness. *Br J Clin Pharmacol*. 2006;61:767-773.
- [186] Al Sayegh F, Almahmeed W, Al Humood S, Marashi M, Bahr A, Al Mahdi H, Bakir S, Al Farhan M. Global Risk Profile Verification in Patients with Venous Thromboembolism (GRIP VTE) in 5 Gulf countries. *Clin Appl ThrombHemost*. 2009 May-Jun;15(3):289-96. doi: 10.1177/1076029608315168.
- [187] Heit JA. Epidemiology of venous thromboembolism. *Nat Rev Cardiol*. 2015; 12:464-474. doi: 10.1038/nrcardio.2015.83.
- [188] Anderson F.A., Wheeler H.B., Goldberg R.J., Hosmer D.W., Patwardhan N.A., Jovanovic B., Forcier A., Dalen J.E. A Population-Based Perspective of the Hospital Incidence and Case-Fatality Rates of Deep Vein Thrombosis and Pulmonary Embolism: The Worcester DVT Study. *Arch. Intern. Med*. 1991;151:933-938. doi: 10.1001/archinte.1991.00400050081016.
- [189] Essien EO, Rali P, Mathai SC. Pulmonary Embolism. *Med Clin North Am*. 2019 May;103(3):549-564. doi: 10.1016/j.mcna.2018.12.013. PMID: 30955521.
- [190] LeVarge B, Wright C, Rodriguez-Lopez J. Surgical management of acute and chronic pulmonary embolism. *Clinical Chest Med* 2018; 39: 659-67.
- [191] Fukuda I, Daitoku K. Surgical embolectomy for acute pulmonary thromboembolism. *Ann Vasc Dis* 2017; 10: 107-14.
- [192] Cheng Y-J, Liu Z-H, Yao F-J, et al. . Current and former smoking and risk for venous thromboembolism: a systematic review and meta-analysis. *PLoS Med*. 2013;10(9):e1001515. doi:10.1371/journal.pmed.1001515.
- [193] Mahmoodi BK, Cushman M, Anne Næss I, et al. Association of traditional cardiovascular risk factors with venous thromboembolism: an individual participant data meta-analysis of prospective studies. *Circulation*. 2017;135(1):7-16. doi:10.1161/CIRCULATIONAHA.116.024507.
- [194] Gregson J, Kaptoge S, Bolton T, Pennells L, et al. Emerging Risk Factors Collaboration. Cardiovascular risk factors associated with venous thromboembolism. *JAMA Cardiol* 2019;4:163-173.
- [195] Al-Nasser B. Influence of Tobacco Smoking on Perioperative Risk of Venous Thromboembolism. *Turk J AnaesthesiolReanim*. 2020;48(1):11-16. doi:10.5152/TJAR.2019.08683.
- [196] Enga KF, Braekkan SK, Hansen-Krone IJ, le Cessie S, Rosendaal FR, Hansen J-B. Cigarette smoking and the risk of venous thromboembolism: the Tromsø Study. *J ThrombHaemost*. 2012;10(10):2068-2074. doi:10.1111/j.1538-7836.2012.04880.x.
- [197] Sweetland S, Parkin L, Balkwill A, Green J, Reeves G, Beral V Million Women Study Collaborators. Smoking, surgery, and venous thromboembolism risk in women: United Kingdom cohort study. *Circulation*. 2013;127:1276-82. doi: 10.1161/CIRCULATIONAHA.113.001428.
- [198] Robertson S, Miller MR. Ambient air pollution and thrombosis. Part I. *FibreToxicol*. 2018;15(1):1.; Mills NL, Donaldson K, hadoke PW. Adverse cardiovascular effects of air pollution. *Nature Clin Pract*. 2009;6(1):36-44.
- [199] Münzel T, Gori T, Al-Kindi S, et al. Effects of gaseous and solid constituents of air pollution on endothelial function. *Eur Heart J*. 2018;39(38):3543-3550.
- [200] Eliasson B. Cigarette smoking and diabetes. *Prog Cardiovasc Dis* 45: 405-413, 2003. doi:10.1016/S0033-0620(03)80004-X.
- [201] Craig WY, Palomaki GE, Haddow JE. Cigarette smoking and serum lipid and lipoprotein concentrations: an analysis of published data. *BMJ* 298: 784-788, 1989. doi:10.1136/bmj.298.6676.784.
- [202] Chioloro A, Faeh D, Paccaud F, Cornuz J. Consequences of smoking for body weight, body fat distribution, and insulin resistance. *Am J Clin Nutr* 87: 801-809, 2008. doi:10.1093/ajcn/87.4.801.
- [203] Oh SS, Jang JE, Lee DW, Park EC, Jang SI. Cigarette type or smoking history: Which has a greater impact on the metabolic syndrome and its components? *Sci Rep*. 2020 Jun 26;10(1):10467. doi: 10.1038/s41598-020-67524-2.
- [204] Wang T, Jiang CQ, Xu L, et al. The mediating role of inflammation in the association between cigarette smoking and intima-media thickness: The Guangzhou biobank cohort study. *Medicine (Baltimore)*. 2020;99(8):e19207. doi:10.1097/MD.00000000000019207
- [205] Pinto-Sietsma SJ, Mulder J, Janssen WM, Hillege HL, de Zeeuw D, de Jong PE. Smoking is related to albuminuria and abnormal renal function in nondiabetic persons. *Ann Intern Med* 133: 585-591, 2000. doi:10.7326/0003-4819-133-8-200010170-00008.
- [206] Volpe M, Battistoni A, Gallo G, Rubattu S, Tocci G, Writing C, et al. Executive summary of the 2018 joint consensus document on Cardiovascular Disease Prevention in Italy. *High Blood Pressure Cardiovasc Prevent Off J Italian Soc Hypertens*. 2018;25(3):327-341.
- [207] Prescott E, Hippe M, Schnohr P, Hein HO, Vestbo J. Smoking, and risk of myocardial infarction in women and men: longitudinal population study. *BMJ*. 1998; 316:1043-1047. doi: 10.1136/bmj.316.7137.1043.
- [208] Thun MJ, Carter BD, Feskanich D, et al. 50-year trends in smoking-related mortality in the United States. *N Engl J Med*. 2013;368(4):351-364. doi:10.1056/NEJMsa1211127.
- [209] Kotseva K, Wood D, De Backer G, De Bacquer D, Pyorala K, Keil U, Group ES EUROASPIRE III: a survey on the lifestyle, risk factors and use of cardioprotective drug therapies in coronary patients from 22 European countries. *Eur J Cardiovasc PrevRehabil*. 2009;16:121-137. doi: 10.1097/HJR.0b013e3283294b1d.