

International Multispeciality Journal of Health

ISSN: 2395-6291



www.imjhealth.org

Volume-9, Issue-3, March 2023

Preface

We would like to present, with great pleasure, the inaugural volume-9, Issue-3, March 2023, of a scholarly journal, *International Multispeciality Journal of Health*. This journal is part of the AD Publications series *in the field of Medical, Health and Pharmaceutical Research Development*, and is devoted to the gamut of Medical, Health and Pharmaceutical issues, from theoretical aspects to application-dependent studies and the validation of emerging technologies.

This journal was envisioned and founded to represent the growing needs of Medical, Health and Pharmaceutical as an emerging and increasingly vital field, now widely recognized as an integral part of scientific and technical statistics investigations. Its mission is to become a voice of the Medical, Health and Pharmaceutical community, addressing researchers and practitioners in below areas

Clinical Specialty and Super-specialty Medical Science:

It includes articles related to General Medicine, General Surgery, Gynecology & Obstetrics, Pediatrics, Anesthesia, Ophthalmology, Orthopedics, Otorhinolaryngology (ENT), Physical Medicine & Rehabilitation, Dermatology & Venereology, Psychiatry, Radio Diagnosis, Cardiology Medicine, Cardiothoracic Surgery, Neurology Medicine, Neurosurgery, Pediatric Surgery, Plastic Surgery, Gastroenterology, Gastrointestinal Surgery, Pulmonary Medicine, Immunology & Immunogenetics, Transfusion Medicine (Blood Bank), Hematology, Biomedical Engineering, Biophysics, Biostatistics, Biotechnology, Health Administration, Health Planning and Management, Hospital Management, Nephrology, Urology, Endocrinology, Reproductive Biology, Radiotherapy, Oncology and Geriatric Medicine.

Para-clinical Medical Science:

It includes articles related to Pathology, Microbiology, Forensic Medicine and Toxicology, Community Medicine and Pharmacology.

Basic Medical Science:

It includes articles related to Anatomy, Physiology and Biochemistry.

Spiritual Health Science:

It includes articles related to Yoga, Meditation, Pranayam and Chakra-healing.

Each article in this issue provides an example of a concrete industrial application or a case study of the presented methodology to amplify the impact of the contribution. We are very thankful to everybody within

that community who supported the idea of creating a new Research with *IMJ Health*. We are certain that this issue will be followed by many others, reporting new developments in the Medical, Health and Pharmaceutical Research Science field. This issue would not have been possible without the great support of the Reviewer, Editorial Board members and also with our Advisory Board Members, and we would like to express our sincere thanks to all of them. We would also like to express our gratitude to the editorial staff of AD Publications, who supported us at every stage of the project. It is our hope that this fine collection of articles will be a valuable resource for *IMJ Health* readers and will stimulate further research into the vibrant area of Medical, Health and Pharmaceutical Research.



Dr. Kusum Gaur
(Chief Editor)



Mr. Mukesh Arora
(Managing Editor)

Board Members

Dr. Kusum Gaur (Editor-in-chief)

Dr. Kusum Gaur working as professor Community Medicine and member of Research Review Board of Sawai Man Singh Medical College, Jaipur (Raj) India.

She has awarded with WHO Fellowship for IEC at Bangkok. She has done management course from NIHFW. She has published and present many research paper in India as well as abroad in the field of community medicine and medical education. She has developed Socio-economic Status Scale (Gaur's SES) and Spiritual Health Assessment Scale (SHAS). She is 1st author of a book entitled " Community Medicine: Practical Guide and Logbook.

Research Area: Community Medicine, Biostatics, Epidemiology, Health and Hospital Management and Spiritual Health.

Mukesh Arora (Managing Editor)

BE (Electronics & Communication), M.Tech (Digital Communication), currently serving as Assistant Professor in the Department of ECE.

Dr. AMER A. TAQA

Dr. AMER A. TAQA is Professor and Head in Dental Basic Science Mosul University, Mosul, IRAQ. He has been registrar of department of Dental Basic Science Mosul University, Mosul, IRAQ. He has published about 100 of research papers and out of that 50 were of international level. He has awarded many times for scientific researches by Government. He has been member of many examination committees and also is a Member in Iraqi Scientific Staff. He has been working as Editor - reviewer in many journals.

Research Area: Dental Science.

Dr. I.D. Gupta

Dr. I. D. Gupta is Professor Psychiatry and working as additional Principal and Dean of student welfare in SMS Medical College, Jaipur.

He is recipient of Prof. Shiv Gautam oration award by Indian Psychiatric Society. He has done training in YMRS at Monte Carlo and BPRS at Singapore. He has been President Indian Psychiatric Society, Rajasthan State Branch. He is author of "Psycho Somatic Disorder" chapter in 1st edition post graduate text book of Psychiatry by Vyas and Ahuja. He has also worked with National Mental Health Programme and has a lot of publication.

Research Area: Community Mental Health, Psycho somatic and liaison Psychiatry.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY

Dr. Anuradha Yadav

Dr. Anuradha Yadav is working as Professor Physiology, SMS Medical College, Jaipur (Rajsthan) India. She is a popular medical teacher and research scholar who had many publications in indexed journals.

Research Area: CVS & CNS physiology, Medical Education and Spiritual Health.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practcal Guide and Logbook".

Research Area: His area of Interest are Epidemiology, Biostatistics and Spiritual Health.

Prof. Dillip Kumar Parida

Professor and Head in the Department of Oncology, AIIMS, Bhubaneswar.

He has done the Professional Training in Japan (Osaka University, NIBI, AHCC Research Association, Hyogo Ion Beam Center), ESTRO Fellowship in Denmark and India(AIIMS Delhi, BARC Mumbai, SCB Medical College-Cuttak, MKCG Medical College-Berhampur).

Research Area: Brachytherapy, Total Skin Electron Irradiation, Palliative Radiotherapy, Stereotactic & Conformal radiotherapy, Radiation Cell Biology, Cancer Genetics.

Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY.

Dr Rajeev Sharma (MS; FMAS; FIMSA;FCLS)

He is working as Professor, Department of Surgery, Government Medical College, Chandigarh, India. He has done FMAS, FIMSA and FCLS along with MS (Gen Surgery).

He has about 50 international and national publications to his credit. He has held various positions in the Association of Minimal Access Surgeons of India (AMASI) from time to time. He has also acted as instructor of various AMASI skill courses held at different places in India. He has established Surgical Technique learning centre at GMCH Chandigarh for imparting training to the budding surgeons in the field of minimal access surgery. He is also the reviewer in the subject in various journals.

Research Area: Minimal Access Surgery.

Dr Anshu Sharma (MS ANATOMY)

She is Presently working as assistant professor in the department of Anatomy, GMCH, Chandigarh. She has many publications in various national and international journals. She is executive member of Anatomical Society of India (ASI) and North Chapter of ASI. She is also a member of editorial board of Journal of Medical College Chandigarh.

Research Area: Congenital Malformation, Developmental Anatomy.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practical Guide and Logbook".

Research Area: His areas of Interest are Epidemiology, Biostatistics and Spiritual Health.

Dr. Dilip Ramlakhyani

Dr. Dilip Ramlakhyani working as Associate professor Pathology and member of IT Committee of Sawai Man Singh Medical College, Jaipur (Raj) India. He has published many articles in indexed journals.

Dr. Virendra Singh

Dr. Virendra Singh worked as Supernatant and head of department of Pulmonary Medicine, SMS Medical College, Jaipur (Rajsthan) India.

He has gone abroad for many training courses and to present research papers. He had been chairman of Research Review Board of SMS Medical College, Jaipur. He is a great research scholar and had published book related to his faculty and had many publications in indexed journals.

Dr. Mahesh Sharma

Dr. Mahesh Sharma is a Principle specialist General Surgery in Rajasthan State Government, India. He has been PMO of district hospitals for more than 15 years. He has gone abroad as observer of many of training related to his speciality. He has published and present many research paper in India as well as abroad.

He has developed Spiritual Health Assessment Scale (SHAS) with Dr. Kusum Gaur.

Research Area: General Surgery, Health and Hospital management and Spiritual Health.

Dr. Ravindra Manohar

Professor Community Medicine, working as head of department of PSM, SMS Medical College, Jaipur (Rajsthan) India.

Previously he has worked in BP Kiorala Institute of Medical Sciences, Nepal. He has visited CDC Atlántica for a Statistical workshop. He has been convener of MBBS and PG exams. He is a research scholar and had many publications in indexed journals.

Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Table of Contents

Volume-9, Issue-3, March 2023

S.No	Title	Page No.
1	Tackling Bipolar Disease with Impulsivity: The Role of Endoxifen Authors: Dhvanit Dave  DOI: https://dx.doi.org/10.5281/zenodo.7785297  Digital Identification Number: IMJH-MAR-2023-1	01-03
2	Knowledge, Attitude and Practice of standard Precautions among Selected Health care Workers in University of Port Hacourt Teaching Hospital, Rivers State Authors: O.E, Mbenyi, E.A. Nwokea, C.O.A. Amadia  DOI: https://dx.doi.org/10.5281/zenodo.7785303  Digital Identification Number: IMJH-MAR-2023-2	04-15

Tackling Bipolar Disease with Impulsivity: The Role of Endoxifen

Dhvanit Dave

Head of Psychiatry, Deep Hospital and Deaddiction centre, Bhuj, Gujarat, India
Nr. Shrushti Park, Commerce College Rd, Bhuj, Gujarat 370001, India

Received:- 02 March 2023/ Revised:- 13 March 2023/ Accepted: 19 March 2023/ Published: 31-03-2023

Copyright @ 2021 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— Bipolar disorder (BD) is characterized by mood changes, and traits of impulsivity are often reported in these patients. Alcohol use disorder is also reported among patients with BD. Protein kinase C is known to be involved in the pathogenesis of BD, and there are reports on its link to impulsivity as well as alcohol use disorder. Thus, the use of a protein kinase C inhibitor, such as endoxifen, could address all three conditions. We present a case of a male patient with BD who displayed impulsivity and alcohol abuse. Initial management with combination therapy was not successful in reducing symptoms. The addition of endoxifen to the treatment allowed the reduction of dosage of sodium valproate and lithium and led to a rapid improvement in symptoms. This report showcases the utility of endoxifen in the management of bipolar disorder.

Keywords— Endoxifen, bipolar disorder, impulsivity, protein kinase C.

I. INTRODUCTION

Bipolar disorder is a chronic mental health disorder characterized by drastic mood changes. Bipolar mania is linked to overactive protein kinase C (PKC) signaling. Patients experience acute manic or mixed episodes, with changes in mood, energy, and activities.¹ Impulsivity is also noted in patients with bipolar disorder during periods of acute mania. This behavior encompasses uncontrolled expenditure and suicide attempts.² Furthermore, there are reports that patients with comorbid bipolar disorder and alcohol use disorders demonstrate greater impulsivity compared with bipolar disorder patients without alcohol misuse. Conversely, impulsivity associated with bipolar disorder may increase the risk for impulsivity-related disorders such as alcohol use disorder.³

We present in this report, a case of a patient with bipolar disorder who showed traits of impulsivity as well as alcohol abuse. The case describes the use of combination therapy and the introduction of endoxifen in the regimen, which led to rapid improvement in bipolar symptoms.

II. CASE REPORT

A 38-year-old male (weight 86 kg) was admitted to the hospital for a sudden-onset seizure. Blood tests were normal, and the patient underwent magnetic resonance imaging (MRI) of the brain with epilepsy protocol, as well as electroencephalography (EEG). On the bases of these tests, the patient was diagnosed with mesial temporal sclerosis (MTS). He was treated with oxcarbazepine 600 mg, brivaracetam 100 mg, and clobazam 10 mg. The patient initially improved with this treatment, but then began feeling low and lost interest in activities he previously found pleasurable. The patient stopped going to work (he was a crane operator at a shipping port) and began drinking alcohol daily (8-10 pegs of whiskey or rum).

Due to his chronic alcoholism, he underwent a major accident and was admitted to the neurosurgical department once again. The patient had suffered undisplaced facial bone fractures and subdural hemorrhage. The neurosurgeon started the patient on antidepressants. The patient showed improvement within 4 weeks. Subsequently, the patient started talking grandiosely and started purchasing unnecessarily and spending money on buying products that were of no use. The patient had purchased a tractor and spent 50,000 rupees in trading.

At this point, the patient was brought to a neuropsychiatrist for further management. During the assessment of history, it was discovered that the patient had been suffering from bipolar mood disorder type 2 for several years. The Young Mania Rating

Scale (YMRS) score was 47. His uncle also suffered the same illness and committed suicide due to defaulting on treatment. The patient was initiated on lithium 900 mg, sodium valproate 2 g, olanzapine 10 mg HS, trifluoperazine 10 mg, oxcarbazepine 600 mg, and brivaracetam 100 mg.

The patient stabilized with these medications, but suffered side effects of sexual dysfunction, hair loss, and mild tremors. Therefore, sodium valproate was tapered to 1,500 mg and lithium was tapered to 600 mg. However, the patient continued to report tremors and irritability, and was hypomanic even with polypharmacy. The patient's wife reported that the manic episode was 17 days old, and the patient was hypomanic for the last 3 months, with a YMRS score of 35. The patient was initiated on endoxifen 8 mg once daily, and responded well to this treatment, with an improvement in the YMRS score from 35 to 29 in just four days of treatment with endoxifen. The patient's relatives reported that he was calm and composed. After 20 days of treatment with endoxifen, the YMRS score reduced to 13, and therefore, dosages of sodium valproate and lithium were tapered; after 6 weeks, the score was 8; and after 3 months, the score was 3 (Figure 1). At this point, the patient was on endoxifen 8 mg, lithium 400 mg, oxcarbazepine 450 mg for MTS, brivaracetam 50 mg, and clonazepam 1 mg. The patient did not have any manic or depressive symptoms, or impulsivity. Endoxifen treatment led to reduced impulsivity and alcoholism, Furthermore, the patient completely stopped alcohol consumption after 3 months, and resumed his job.




FIGURE 1: Change in Young Mania Rating Scale after initiating treatment with endoxifen

III. DISCUSSION

The case described above demonstrates how switching to endoxifen allowed a reduction of the dosages of other drugs (sodium valproate and lithium) and led to a rapid improvement of symptoms. The treatment with endoxifen was well-tolerated, and the patient was able to resume his job as well as stop alcohol consumption. Endoxifen proved to be a useful part of this combination treatment regimen in the case of a young male with comorbid bipolar disorder and alcohol abuse.

Endoxifen is a metabolite of tamoxifen and has a fourfold higher inhibitory action against PKC compared to the parent molecule. Endoxifen has a number of advantages, including its lipid-soluble nature and ability to cross the blood-brain barrier.¹ PKC plays a role in the pathogenesis of not just bipolar disorder, but also impulsivity and alcohol use disorder.⁴⁻⁶ Patients with mood disorders have an imbalance in PKC signaling, which is required for the modulation of neuronal processes involved in mood regulation.⁴ Studies have shown that elevated levels of PKC signaling can contribute to impulsivity, which is a sign of prefrontal cortical dysfunction leading to impulsivity.⁵ Finally, PKC is involved in the response to ethanol exposure, wherein animal studies show that deletion of PKC produces a low ethanol consumption.⁶ These studies indicate that targeting PKC may be a useful approach to treating these conditions.^{4,6}

Clinical trials have demonstrated the safety and effectiveness of endoxifen, a PKC inhibitor, for the management of bipolar disorder. At a dose of 8 mg daily, endoxifen significantly reduced YMRS by 21 days of treatment (from 33.1 to 17.8).¹ In the case presented in this report as well, the YMRS score reduced from 35 to 13 after 20 days of treatment with endoxifen. Clinical

studies report no serious adverse effects of endoxifen.¹ Similarly, another study reported a reduction of YMRS score by 16.21 points after 21 days of treatment with endoxifen, while it has a safety comparable with that of divalproex.⁴

The patient in this case was a chronic alcoholic, and alcohol can aggravate the course of bipolar disorder, as well as change the expression of PKC. Endoxifen was useful in relieving bipolar disorder symptoms and could also have allowed the de-addiction from alcohol. Past case reports indicate that endoxifen may have a role to play in patients with bipolar disorder and comorbid alcohol abuse.⁷

Clinical trials have reported an early response after four days of treatment with endoxifen, with significant improvement in YMRS score.^{1,4} Our case as well showed a reduction in YMRS score from 35 to 29 in four days. The rapid action of endoxifen is beneficial in providing relief to patients. The lower doses required and lack of interaction with other drugs gives endoxifen a unique advantage over other molecules.^{1,4}

IV. CONCLUSION

In conclusion, endoxifen is an effective treatment option for bipolar disorder and impulsivity, and it reduces the overall burden of other medications that have to be prescribed to such patients. In this case, endoxifen treatment enabled rapid control of symptoms within 4 days, and the patient subsequently returned to work. The patient did not resume alcohol consumption while on endoxifen treatment. Endoxifen is effective as well as safe and can be part of a combination treatment regimen.

REFERENCES

- [1] Ahmed A, Sheikh S, Khan MA, Chaturvedi A, Patel P, Patel R, et al. (2021). Endoxifen: A new, protein kinase C inhibitor to treat acute and mixed mania associated with bipolar I disorder. *Bipolar Disorders*. 23(6): 595–603. <https://onlinelibrary.wiley.com/doi/10.1111/bdi.13041>.
- [2] Powers RL, Russo M, Mahon K, Brand J, Braga RJ, Malhotra AM, et al. (2013). Impulsivity in bipolar disorder: Relationships with neurocognitive dysfunction and substance use history. *Bipolar Disorders*. 15(8): 876–84. <https://onlinelibrary.wiley.com/doi/abs/10.1111/bdi.12124>.
- [3] Jen A, Saunders EFH, Ornstein RM, Kamali M, McInnis MG. (2013). Impulsivity, anxiety, and alcohol misuse in bipolar disorder comorbid with eating disorders. *International Journal of Bipolar Disorders*. 1: 13. <https://journalbipolar disorders.springeropen.com/articles/10.1186/2194-7511-1-13>.
- [4] Ahmed A, Sheikh S, Reddy MS, et al. (2016). Endoxifen, a new treatment option for mania: A double-blind, active-controlled trial demonstrates the antimanic efficacy of endoxifen. *Clinical and Translational Science*. 9(5): 252–9. <https://ascpt.onlinelibrary.wiley.com/doi/10.1111/cts.12407>.
- [5] Birnbaum SG, Yuan PX, Wang M, Vijayraghavan S, Bloom AK, Davis DJ, et al. (2004). Protein kinase C overactivity impairs prefrontal cortical regulation of working memory. *Science*. 306(5697): 882–4. <https://www.science.org/doi/10.1126/science.1100021>.
- [6] Newton PM, Ron D. (2007). Protein kinase C and alcohol addiction. *Pharmacological Research*. 55(6): 570–7. [https://linkinghub.elsevier.com/retrieve/pii/S1043-6618\(07\)00087-4](https://linkinghub.elsevier.com/retrieve/pii/S1043-6618(07)00087-4).
- [7] Dubey V. (2022). Long-term effectiveness and safety of endoxifen in the treatment of bipolar mania: A case report. *International Journal of Advances in Medicine*. 9(2): 169–72. <https://www.ijmedicine.com/index.php/ijam/article/view/3311>.

Knowledge, Attitude and Practice of standard Precautions among Selected Health care Workers in University of Port Harcourt Teaching Hospital, Rivers State

O.E, Mbenyi^{1*}, E.A. Nwokea², C.O.A. Amadia³

Department of Public Health, School of Health Technology, Owerri, Imo State, Nigeria.

*Corresponding author

Received:- 07 March 2023/ Revised:- 13 March 2023/ Accepted: 20 March 2023/ Published: 31-03-2023

Copyright @ 2021 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— This research study centered on determining the knowledge, attitude, and practice of standard precautions among selected healthcare workers in the University Teaching Hospital, Port Harcourt, Rivers State. The specific objectives are to examine the socio-demographic characteristics of the respondents, assess the level of awareness of standard precautions among healthcare workers, ascertain the attitude of healthcare workers in the University of Port Harcourt Teaching Hospital, Rivers State towards standard precautions, and determine the level of practice of standard precaution. Related literature reviewed. A descriptive survey design was employed in this study. Out of the 400 health workers, Taro Yamane's sample size determination gave a sample size of 200. A well-structured questionnaire was used to extract information from the respondents. Descriptive statistics such as means, and frequency distribution and percentages were used to analyze all the objectives. Chi-Square was used to analyze the hypothesis posed to the study. The result revealed that there were more females (67.5%) who participated in the studies than their male counterparts (32.5%). The majority (42.0%) of the respondents were still within their mid-age of 30 – 39 years, followed by 26% who were within the age range of 40 – 49 years of age. Furthermore, the study showed that all participants (100%) had at one time or the other previous knowledge of standard precaution as a concept in the health system. However, their sources of information varied from formal training (65%), colleague/friend (26%) media (5.5), and other means (3.5%). Also, the respondents opined that every employer in the health system should as a matter of utmost importance make provision for training on standard precautions. They believed that standard precaution is helpful in the protection against health hazards in the workplace (= 3.40). Some (50%) thought that recap of the needle after use is a safe way to prevent needle injury and transmission of diseases. Safety measures agreed upon by the respondents 98% include disposal of needles and syringes immediately after use. Method of needle disposal in UPTH include the use of waste bin (50.5%) and enclose and burn (60%). The Chi-Square value (9.88) and p-value ($p > 0.05$) means that we do not have sufficient evidence to reject the null hypothesis which states that there is no significant impact of knowledge of standard precaution on practice.

Keywords— Port Harcourt, health care workers, standard precautions, blood, attitude.

I. INTRODUCTION

A set of guidelines aimed at protecting health care workers (HCWs) from blood-borne infections is known as universal precautions (1). The concept of "universal precautions" was first proposed by the Centers for Disease Control and Prevention (CDC) in 1987 to protect health care workers from exposure to blood-borne pathogens. The Centers for Disease Control and Prevention (CDC) recommended in 1996 that universal precautions be renamed standard precautions, which combine the major features of universal precautions and body substance isolation (BSI). The precautions apply to all body fluids, including blood, secretions, and excretions (except sweat), regardless of whether they contain visible blood, skin that is not intact, mucous membranes, or any unfixed tissue or organ (other than intact skin) from a human (living or dead), HIV, or hepatitis containing culture medium or other solutions. A successful Infection Prevention and Control (IPAC) plan must be initiated and implemented in any healthcare facility before an evaluation of the knowledge, attitude, and practice (KAP) of standard precautions by healthcare personnel can be conducted. According to their professional group and level of experience, among other things, HCW exhibit variable KAP of standard precautions, according to numerous studies. Improved standard precaution

compliance has been linked to increased professional experience, standard precaution knowledge and training, and high-risk perception among health professionals. The majority of studies from around the world have shown that healthcare workers are more likely to follow standard precautions (2), but some others have not (3).

Despite research on standard precautions compliance among Nigerian healthcare professionals (4), the Nigerian community lacks a clear understanding of the professional disparities in KAP of standard precautions of infection control among healthcare professionals. So, our goal was to look at how well doctors, nurses, and laboratory personnel at two tertiary hospitals in Nigeria were following some of the basic measures. In order to ascertain the knowledge, attitudes, and application of conventional precautions among health professionals at the university teaching hospital in Port Harcourt, Rivers State—a developing metropolitan area with rising HIV/AIDS and other infectious disease cases—this study was created.

1.1 Problem statement:

- As they carry out their clinical duties in the hospital, healthcare professionals (HCWs) run the danger of occupational risks. Roughly 12% of the working population, or about 35 million individuals, work in the health care industry, but their occupational health has been neglected (WHO,2007). Sharps injuries and contact with deep body fluids expose them to blood-borne pathogens like HIV, hepatitis B, and hepatitis C viruses.
- The Occupational Safety and Health Administration estimates that 5.6 million HCWs worldwide who handle sharp devices, are at risk of occupational exposure to blood borne pathogens. Often, these injuries go unreported for a variety of reasons, including stigma. The total HIV sero-prevalence was 4.4% according to the Nigerian sero-sentinel survey from 2005. The high incidence in the nation puts HCWs at risk for occupational disease (3).
- Inadequate sharps containers and disposal facilities, bad practices like bending needles, recapping needles, detaching needles, reusing needles, a lack of supply of injection equipment, and unwarranted and unsafe injection practices, which put both patients and healthcare workers at risk, may all contribute to the rising prevalence of morbidity and mortality following exposure to blood-borne infections. In light of the aforementioned issues, the researcher made the decision to conduct a study on HCWs' knowledge, attitudes, and practices regarding standard precaution at the university of Portharcourt teaching hospital in the state of rivers

1.2 Objective of the study

1.2.1 General objectives

The general objective of this research is to determine the knowledge, attitude and practice of standard precautions among health care workers in the University Teaching hospital, Portharcourt, vers State.

1.2.2 Specific objectives

The research objectives are

- a) To assess the level of awareness of standard precautions among health care workers in University of Port Harcourt Teaching hospital, Rivers State.
- b) To ascertain the attitude of health care workers in University of Port Harcourt Teaching hospital, Rivers State towards standard precautions.
- c) To determine the level of practice of standard precautions among health care workers in public secondary health facilities in University of Port Harcourt Teaching hospital, Rivers State.

II. METHODS

The study was carried out at the University of Port harcourt Teaching Hospital (UPTH) Rivers State, Nigeria. Descriptive survey method was utilized in this investigation. A descriptive survey study aims to give information on a situation's nature and status as it is at the time of the study.

UPTH policy on standard precaution and procedure on universal precautions and sanitary policy practices.

2.1 Purpose

This policy's objectives are infection control, communicable disease prevention, and the establishment of rules for universal safety precautions and hygienic behaviors, such as hand washing.

2.2 Policy

The UPTH has a policy of using and practising good hygiene. All UPTH employees were get training on hygienic practices, general infection control methods, and universal safeguards to stop the transmission of bloodborne infections. This includes proactive measures to reduce the possibility of catching an infection or disease from another person or from contact with a contaminated surface. Blood, bodily fluids that may clearly be seen to have blood contamination, semen, and vaginal secretions are all infectious materials that should be handled with universal caution. All employees adhered to standardized safety measures and hygienic standards, including:

- 1) Use of appropriate hand washing.
- 2) Putting on gloves when handling contagious materials
- 3) When exposed to potentially dirty or infectious objects, wear a gown or an apron.
- 4) If splashing is likely, wear a mask and goggles.
- 5) Put on gloves and wipe affected surfaces with disinfectant
- 6) Handling sharps properly
- 7) Put on gloves and wash contaminated laundry with correct bagging techniques.

III. PROCEDURE

Universal Precautions and Infection Prevention and Control.

1. The idea of universal precautions assumes that all bodily fluids are infectious, hence all medical professionals must handle blood and/or bodily fluids as though they are infected. This not only safeguards caregivers but also eliminates any stigma connected with using gloves and masks around patients who are very contagious by following the same protocols for everyone.
2. The single most crucial behavior for halting the spread of infection and disease is hand washing. Regardless of the existence or absence of any known diseases and infections, proper hand washing were performed as part of regular work practices and routines. Additionally, staff members are expected to help customers wash their hands frequently. Every employee washed their hands at the start and conclusion of every shift, before and after handling food, before and after helping someone with personal care, after any contact with blood or bodily fluids, and after any potential contamination. Examples include managing dirty clothing or dishes, taking out the trash, etc. The following procedure were followed when washing hands.
 - a) Use warm water on your hands and wrists.
 - b) Use sufficient soap to produce lather.
 - c) Make lather by rubbing your hands together.
 - d) Lather up your hands and vigorously rub them for at least 10 to 15 seconds (palms backs of hands, fingers, spaces between fingers, around and under nails, and wrists). Wash your hands for three to five minutes if you have come into contact with blood or other potentially infectious items.
 - e) Completely rinse with warm water.
 - f) Using a fresh paper towel, pat the skin dry.
 - g) Use a dry paper towel to turn off the faucet.

- h) Get rid of trash.
3. Employees were made sure that their coughs and sneezes were properly covered. Coughing or sneezing into a tissue or paper towel was considered to be appropriately covered. Staff members were cough or sneeze into their elbows if these things were not accessible. Additionally, staff members were expected to help clients understand and employ proper cough- and sneeze-coverage techniques.
 4. Gloves were worn to create a barrier between hands and any potential infection sources. When contact with high risk bodily fluids is reasonably anticipated, gloves must be used. For each circumstance and each individual serviced, new gloves were utilized. To ensure safety, gloves were removed from "skin to skin" and "glove to glove".

3.1 Population For the study

Doctors, nurses, medical laboratory scientists, health aides, and waste handlers at the teaching hospital run by the University of Port Harcourt made up the study's population. There were 400 people working as health care providers who have been employed for at least 6 months and are actively involved in handling garbage, needles, sharp items, and bodily fluids.

3.2 Sample and Sampling technique

3.2.1 Sample size determination:

Using Taro Yamane formula for the sample size.

$$n = \frac{N}{1 + N(e)^2}$$

Where, N = population of study

n= sample size

e= level of significance or margin of error

I= unit (a constant)

Therefore,

$$N = 400$$

$$e = 0.05$$

$$n = ?$$

(The choice of 0.05 level of significance is purely an exclusive decision of researcher).

Replacing the values above with the above formulae, we have

$$n = \frac{400}{1 + 400(0.05)^2}$$

$$n = \frac{400}{1 + 400(0.0025)}$$

$$n = \frac{400}{1 + 1}$$

$$n = \frac{400}{2}$$

$$n = 200$$

3.2.2 Sampling method

Multistage sampling technique was used. First stage was the sampling of 30% of the departments in the university of Portharcourt teaching hospital which should be representative of all the twenty departments. 30% of it was 6 departments.

Second stage was purposeful selection of the six departments who were more involved in handling patient blood, bodily fluids, sharp objects, needles and waste. Who also have been employed for at least 6 months. The selected departments were Male and Female medical wards, Male and Female surgical wards, Theatres, medical laboratories, obstetrics and Waste handlers.

Third stage is the sampling of the health care workers from the selected departments where they work using simple random sampling. 14 doctors, 19 nurses, and 11 health assistants were randomly sampled from male and female medical wards. Same number respectively sample from male and female surgical wards. In the theatre, 12 doctors, 18 nurses, and 11 health assistants were sample. From the laboratories 33 medical laboratory scientists and 11 health assistants sampled. Also 27 waste handlers were sampled in waste handling department.

3.2.3 Instrument of Data Collection

The descriptive survey research employs the use of structured questionnaires, which was thought to be the most effective method for reaching the study's population, particularly when the instrument can be used to collect the necessary data. A questionnaire with both open-ended and closed-ended questions was the research tool employed for the study. Based on the research topics put forth for the study, the questionnaire was created from the review of the literature. The questionnaire's Section "A" contained information about the respondents' backgrounds. Section B investigated standard precaution knowledge, Section C studied standard precaution attitudes, and Section D examined standard precaution practices.

3.2.3.1 Validity of the Instrument

The characteristics of tests and measurements are what are referred to as validity. Validity is the degree to which a measuring device measures what it is designed to measure. The researcher created the questionnaire as the tool for gathering data, and the project supervisor reviewed it critically before making suggestions. Two experts validated the updated version of the survey for relevance to the objectives, material coverage, and language usage.

3.2.3.2 Reliability of the Instrument

Using the test-retest procedure, the instrument's dependability was evaluated. In order to conduct the test, copies of the questionnaire were distributed to twenty (20) respondents who were **No table of contents entries found**. Representative of the target demographic. The same procedure was repeated a week later in order to evaluate the veracity of the respondents' responses. The Crombach Alpha test was used to determine whether the results were consistent, and a coefficient of 0.7 was obtained, indicating that the instrument was deemed to be dependable.

3.2.4 Method of Data Collection

The respondents were given instructions on how to complete the questionnaires as well as the study's goal. Data collection for the project took a few weeks, and during that time I made myself available to respond to questions that were loosely related to it. The questionnaire was used to gather data. After receiving informed consent, the questionnaire was given to the respondents at each department. .copies of the questionnaire that have been completed, collected on the spot.

3.2.5 Method of Data Analysis

Data were collected, entered into the computer and analyzed using SPSS Version 21(Statistical Package for Social Science) In order to analyze the data, descriptive statistics (frequencies, tabular percentages, mean score and standard deviation), as well as the chi-square test, were utilized. **Ethical considerations/Informed consent.**

The Federal University of Technology, Owerri, School of Health Technology Ethical Review Board granted ethical approval. Before giving out the questionnaires', informed consent was obtained from the respondents.

IV. RESULTS

4.1 Socio-Demographic Characteristics of The Respondents in UPTH

TABLE 1a
SOCIO-DEMOGRAPHIC DISTRIBUTION OF HEALTH WORKERS (N=200)

Demographic variables	Frequency	Percentages (%)
Gender		
Female	135	67.5
Male	65	32.5
Total	200	100
Age range		
20-29	45	22.5
30 – 39	84	42.0
40 – 49	52	26.0
50 – above	19	9.5
Total	200	100
Years of service		
1 – 10	116	58.0
11 – 20	45	22.5
21 – 30	36	18.0
31 – 35	3	1.5
Total	200	100

The sociodemographic distribution of health workers in UPTH is shown in table 1a above. The findings showed that 67.5% of the participants were female, as opposed to 32.5% of the male peers. The majority (42.0%) of respondents were between the ages of 30 and 39, with 26% of respondents being between the ages of 40 and 49. 9.5% of the population is over 50 and represents a tiny percentage of the older population. Additionally, the bulk of respondents were still in their first one to ten years of employment, with another 22.5% staying for eleven to twenty years.

TABLE 1b
SOCIO-DEMOGRAPHIC DISTRIBUTION OF HEALTH WORKERS (N=200)

Demographic variables	Frequency	Percentages
Marital status		
Married	114	57.0
Single	62	31.0
Widow/widower	13	6.5
Divorced/separated	11	5.5
Religion		
Christianity	177	88.5
Islam	14	7.0
Traditional	9	4.5
Total	200	100
Occupation		
Nursing	56	8.0
Doctor	40	20.0
Laboratory Sci.	33	16.5
Waste handlers	27	13.5
Health assistants	44	22
Total	200	100

Results of analyses on marital status, religion, and occupation are shown in Table 1b above. The majority (57.0%) of the responding health professionals were married; 31% were single, and the remaining 12% had marital issues ranging from separation to the loss of a spouse. The majority of respondents (85.5%) practiced Christianity, 7% practiced Islam, and only 4.5% stated traditional religion. In this activity, nurses make up the majority (28%) of participants, followed by health assistants (22%) and physicians of medicine (20%). Lab scientists (16.5%) and Waste handlers (13.5%) round out the list.

4.2 Level of Awareness (Knowledge) of the Respondents in UPTH On Standard Precaution

TABLE 2a
KNOWLEDGE AND SOURCE OF INFORMATION ON STANDARD PRECAUTION

Demographic variables	Frequency	Percentages	SA	A	Indifferent	SD	Mean
Awareness							
Previous knowledge of standard precaution as a concept in the health system	200	100					
Source of information							
Formal training	130	65.0					
Colleague/friend	52	26.0					
Media	11	5.5					
Others	7	3.5					
Total	200	100					
Components of standard precaution							
Hand hygiene			84	11	-		3.42
Use of personal protective equipment			128	72	-		3.64
Safe injection practices			72	10	20		3.26
Safe handling of potentially contaminated equipment surfaces		126	74	-		3.63	
Respiratory hygiene etiquette			81	10	10		3.36
Anal/perineal hygiene			44	13	17		3.14

Mean response > Likert mean (2.5) = significant

TABLE 2b
KNOWLEDGE AND SOURCE OF INFORMATION ON STANDARD PRECAUTION

Items	Strongly Agree (SA)	Agree (A)	Indifferent	Mean
Advantages of standard precaution				
Protects both health workers and patients	121	74	5	3.56
Reduced spread of communicable disease	48	93	49	2.85
Not associated with stigma and discrimination	56	122	22	3.17

Results on the respondents' knowledge of standard precaution and the sources from which they learned about it are shown in Tables 2a and 2b. The investigation reveals that everyone (100%) had some prior understanding of conventional precaution as a concept in the healthcare system. Their informational sources, however, ranged from formal education (65%), colleagues/friends (26%) to the media (5.5) and other sources (3.5%). There is strong evidence that standard precaution knowledge and practices are subjected to deliberate training for health professionals.

The respondents agreed very well that components of standard precaution include hand hygiene ($\bar{x} = 3.42$), use of personal protective equipment (PPE) ($\bar{x} = 3.64$), safe injection practices ($\bar{x} = 3.26$), safe handling of potentially contaminated equipment surfaces ($\bar{x} = 3.63$), respiratory hygiene etiquette ($\bar{x} = 3.36$) and anal/perineal hygiene ($\bar{x} = 3.14$). The response means of items raised under the advantages of standard precaution also revealed that the respondents agreed that practice of

standard precaution has immense benefit of which some of it include protection of both health workers and patients ($\bar{x} = 3.56$), reduced spread of communicable disease ($\bar{x} = 2.85$), dealing with the issue of stigmatization and discrimination ($\bar{x} = 3.17$).

4.3 Attitude of Medical Workers in UPTH towards Standard Precaution

TABLE 3a
ATTITUDE OF THE RESPONDENTS TOWARDS STANDARD PRECAUTION

Variable	Strongly agree	Agree	Indifferent	Disagree	Mean
Attitude towards standard precaution					
Employers should always provide training on standard precautions	27	172	1		3.14
Standard precaution are useful in protecting against hazards in workplace	86	107	7		3.40
Standard precaution are not really necessary in hospitals	-	2	16	182	1.1
They are only necessary for theatre workers	-	28	72	100	1.64
Standard precaution can prevent spread of infections so should be observed always	58	93	49		2.8
Standard precaution requires one to recap needles after use to avoid needle injury and infection	80	120			3.4

n = 200, mean response > 2.5 = significant

The study of the respondents' attitudes about typical precaution techniques was displayed in Table 3a. The respondents believed that it was crucial for every employer in the healthcare system to provide training on common precautions, as evidenced by the mean response of 3.14, which was greater than the likert mean (cut-off). They believed that standard precaution is helpful in the protection against health hazards in workplace ($\bar{x} = 3.40$). The low response mean of 1.1 and 1.6 is an indication that the health workers in UPTH do not believe standard precaution is not really **Table 3b: Attitude of the respondents towards standard precaution n (%)** necessary or that it is mainly for those working in the theatre respectively. On the contrary they believed that standard precaution helps to prevent infections ($\bar{x} = 2.80$) and therefore should always be observed while carrying out any form activity in the hospital.

TABLE 3b
ATTITUDE OF THE RESPONDENTS TOWARDS STANDARD PRECAUTION N (%)

Items	Strongly agree	Agree	Indifferent	Disagree
Management of HIV and HBV positive patients: the following precautionary measures should be adopted while handling patients with the above viruses:				
Wearing double hand gloves	178(89.0)	22(11.0)		
Wearing goggles/face mask	67(33.5)	130(65.0)	3(1.5)	
Wearing gowns/aprons	180(90.0)	20(10.0)		
Wearing boot/wear	170(85.0)	27(13.5)	3(1.5)	

Similar to table 3a, respondents in table 3b responded that it is essential to use double hand gloves (89%), goggles or a face mask (65%), a gown (90%), and boots (85%) when working with patients who have HIV or HBV. The aforementioned evidence amply demonstrates the positive attitude the UPTH personnel has toward routine precaution.

4.4 Practice of Standard Precaution

TABLE 4
PRACTICE OF STANDARD PRECAUTION BY THE RESPONDENTS

Items	Strongly agree	Agree	Indifferent	Disagree
Ways to handle spill of blood and body fluid. They among others include:				
Clean the floor during and after any medical activity.	109(54.5)	91(45.5)		
Clean with an antiseptic immediately	58(29.0)	88(44.0)	54(27.0)	
Use soap and water to clean affected surface	192(96.0)	8(4.0)		
Use 0.5% chlorine bleach	67(33.5)	94(47.0)	39(19.5)	
Injection safety practices				
Recap needle after use	100(50)	66(33.0)	34(17.0)	
Detach needles from syringe immediately after use	-	56(28.0)	69(34.5)	75(37.5)
Dispose needle and syringe immediately after use	196(98.0)	4(2.0)		
Methods of needle disposal				
Use of waste bin	90(45.0)	101(50.5)	9(4.5)	
Enclose and burn	78(39.0)	120(60)	2(1)	
Bury them underground	9(4.5)	37(18.5)	34(17.0)	120(60)
Methods of handling re-usable instruments				
Boil beyond 100°C	77(38.5)	89(44.5)	13(6.5)	21(10.5)
Soak in a disinfectant after washing with soap and water	121(60.5)	66(33.0)	13(6.5)	
Putting in an autoclave	187(93.5)	9(4.5)	4(2.0)	
Sterilize before and after use	188(94)	8(4.0)	4(2.0)	

Results on respondents' use of standard precaution are shown in Table 4. The outcome showed that the personnel had adopted some procedures for dealing blood and bodily fluid spills. The majority (54.5%) clean their floors before and after medical procedures. 44% of people promptly wipe infected goods with an antiseptic. In order to maintain the hospital rooms free of infections, the medical staff at University of Port Harcourt Teaching Hospital also utilizes soap and water to clean infected surfaces (96%) and 0.5% chlorine bleach (94%) were also some ways to disinfect rooms.

Some (50%) thought that recap of needle after use is a safe way to prevent needle injury and transmission of diseases. Safety measures agreed by the respondents include disposal of needles and syringes immediately after use (98%). Method of needle disposal in UPTH include the use of waste bin (50.5%) and enclose and burn (60%).

4.5 Test of hypothesis

Statement of hypothesis: knowledge of standard precautions among health care workers do not significantly influence practices in University of Port Harcourt Teaching hospital, Rivers State.

TABLE 5a
CHI-SQUARE HYPOTHESIS ONE

n=200		Practice of standard precaution			Total
		D	A	SA	
Knowledge of standard precaution	D	0(0)	0(0)	11(3.14)	11
	A	0(0)	59(46.73)	46(29.93)	105
	SA	54(22.68)	30(37.38)	0(0)	84
Total		54	89	57	200

chi square is given by:

$$\chi^2 = \sum \frac{(O-E)^2}{E}$$

Chi square = $43.25 + 3.22 + 1.46 + 19.67 + 8.63 = 76.23$

Table value = 9.488

Since the calculated value is higher than the table value, we have sufficient evidence to reject the null hypothesis

TABLE 5b
CHI-SQUARE OF HYPOTHESIS TWO

		Attitude to standard precaution			Total
		D	A	SA	
Knowledge of standard precaution	A	0(0)	107(62.06)	9(51.62)	116
	SA	7(2.94)	0(0)	77(36.12)	84
Total		7	107	86	200

chi square is given by:

$$\chi^2 = \sum \frac{(O-E)^2}{E}$$

Chi square (χ^2) = $5.66 + 32.54 + 35.18 + 46.42 = 199.80$

Table value = 10.597

Since the calculated value is higher than the table value, we have sufficient evidence to reject the null hypothesis

V. DISCUSSION OF FINDINGS

According to the study, more women (67.5%) than men (32.5%) participated in the studies. These women were mostly between the ages of 40 and 49 and had between one and ten years of work experience. Additionally, the analysis revealed that there were more nurses than doctors. This finding is comparable to that of the study (5) conducted, which found that most health workers were nurses working in the wards. This claim is plain to see given the fact that more nurses were employed to help doctors and other healthcare professionals care for the increasing number of people who were admitted in hospitals. This study is comparable to one that was done in France where nurses made up 44% of the group of HCWs analyzed, and another one that was done in Nigeria where nurses made up 50.7% of all the hospital workers studied. This looks to be an indication of the proportion of nurses among HCWs in general (6).

Given how long ago the policy on SP was created, it is not surprising that there was a high level of awareness about Standard Precaution (SP). None of the health care professionals claimed to be unaware that safety measures should be used throughout any type of medical procedure. Everybody (100%) agreed that they had heard of the idea at some point, whether it was through training (school or a workshop), a friend or coworker, the media, or another source. Additionally, it relates to research conducted in the United Arab Emirates, where 97% of respondents were conversant in the idea (7). (8) both claimed to possess a comparable level of knowledge. This demonstrated the level of interest of HCWs in SP, which is most likely related to their belief that Standard precaution is essential for infection control in contexts involving the public health. The majority of the respondents 82.5%) (in the current survey had favorable attitudes toward universal precaution, which is consistent with other findings (9).

Once more, the demonstrated favorable attitude may be the result of their trust in standard precaution (SP), as the SP policy is a solid package intended to stop the spread of nosocomial diseases and guarantee the safety of HCWs while at work. Wearing of PPE was optimally practiced, which is positive because it is essential for infection management in addition to being a key element of SP. This is consistent with a prior study (10). It might be assumed that the respondents' great familiarity with standard precautions prompted them to view them favorably.

In this survey, the respondents (82.5%) have a good attitude toward wearing gloves, a mask, aprons, and boots although in another study, only 34% of respondents reported wearing gloves and coveralls on a regular basis. The primary explanations for this poor PPE use included sporadic access to PPEs, a lack of time to wear them, and the fact that doing so interferes with clinical duties (11). This demonstrated the need for management and HCWs to work together as closely as possible to implement Standard precaution completely. Secondly, the management should establish an enabling atmosphere by routinely providing

PPEs and other necessary resources and supervising their use, while the HCWs should guarantee complete compliance by adhering to established protocols and constantly utilizing resources.

This study shows that standard precaution is highly practiced in UPTH. Over 50% of the respondents asserted that whenever there is spill of blood or body fluid, such is cleaned immediately with either of antiseptic, use of soap and water or 0.5% chlorine bleach.

Majority (60%) do not detach needles from syringe after use. This is a better standard precaution as detaching and capping after use may dispose a health worker to infections. This practice is in line with research conducted by (12) where respondents some of the respondents confirmed that they do not recap needle after use and detach needles from syringe after use. (13) in their work; assessment of knowledge and practice of standard precautions among nurses working at Federal Medical Centre Gombe also opined that trying to detach or recap needles is not a good practice as it is dangerous and may lead to body injury

The hypothesis test in the work showed that the influence of knowledge on practice is not significant enough 9.88, Since the calculated value is higher than the table value, we have sufficient evidence to reject the null hypothesis. . This may imply that there are additional potentially confounding factors of the use of standard precautions in addition to knowledge or training in infection control. Lack of resources and facilities for infection prevention (IPAC) and control has been reported as a major factor influencing poor practice of IPAC in health facilities in Nigeria (14; 2) and other countries of the world (15). It could also be that our respondents practice standard precaution based on availability of resources and not necessarily based on knowledge. Some health workers knowledgeable but may find it inconveniencing using some safety materials. Excess work and time constraints have also been reported by other studies (14; 16).

VI. CONCLUSION

There has always been a problem with basic precautionary measures, particularly in the Nigerian system. Either there aren't enough safety measures in place, or the medical staff lack the self-control to always use them. This study was carried out to ascertain the level of knowledge, attitude, and standard precaution practice in UPTH. The evidence demonstrated that the medical staff was not at all ignorant of common safety measures. Most of them had formerly taken part in training that taught them the value of following SP standards in order to protect themselves or others. Some people were informed about it through a colleague, media outlet, etc.

While principles of standard precaution should be strictly adhered by surgeons and physicians for their own safety as well as setting leadership roles for their residents, nurses, success in the implementation of SP guidelines depends on so factors such as optimal awareness and a positive attitude in all health care staffs, these goals are not reached without qualified employee receiving continuing education. This study revealed that the health care workers in UPTH has positive attitude towards standard precaution. Bodily fluid or blood spillage were being cleaned immediately they occur with disinfectants to avoid possible infectious contamination. Use of personal protective equipment such as gloves and face masks, and injection safety were some of the safety measures taken by respondents.

Frequent cleaning with antiseptic is been practiced in the hospital. Soap and water were been used for hand hygiene, sometimes chlorine bleach is used to disinfect contaminated surfaces. Even though some healthcare professionals are unawares that recapping needle can be injurious, however needle and syringe were being exposed immediately after use. Method of disposal include use of waste bin, incinerator burning and burying in the ground. In all, all the objective set out for this work were realized.

RECOMMENDATIONS

In order to promote good standard precaution and mitigate the risk of hospital acquired infections, it is necessary for health authorities in Nigeria to institute policies that make it compulsory to establish IPAC (infection control and prevention) committees in all hospitals. Such policies should also outline measures that ensure that IPAC resources are made routinely available and that knowledge and practice of standard precautions are improved through regular IPAC training of hospital staff, with special emphasis on newly qualified health workers such as house officers and staff nurses.

ACKNOWLEDGEMENTS

The authors wish to thank the study participants for their contribution to the research, as well as current and past investigators and staff.

CONFLICTS OF INTEREST

The Author(s) declare(s) that there is no conflict of interest.

Financial support and sponsorship: Nil

REFERENCES

- [1] Bennett, G and Mansell, I. (2014). Universal precautions: a survey of community nurses' experience and practice. *International Journal of Clinical Nursing*. 1(2): 20 – 27.
- [2] Okechukwu, EF., and Modteshi, C. (2012). Knowledge and practice of standard precautions in public health facilities in Abuja, Nigeria. *International Journal of Infection Control* 8: 1–7.
- [3] Shuper, PA., Kiene ,SM., Mahlase, G., Macdonald, S., Christie ,S., Comman, D. H., Fisher, W. A., Greener, R., Lalloo, U. G., Pillay, S., van Loggelenberg, F., and Fisher, JD. (2014). HIV transmission risk behavior among hiv-positive patients receiving antiretroviral therapy in KwaZulu-Natal, South Africa. *AIDS Behaviour* 18: 1532–1540.
- [4] Ofili ,AN., Asuzu, MC., and Okojie, OH. (2008). Knowledge and practice of universal precautions among nurses in central hospital, Benin-City, Edo State, Nigeria. *Nigerian Postgraduate Medical Journal* 10: 26–31.
- [5] Arinze- Onyia SU, Ndu AC, Aguwa EN, Modebe I, Nwamoh UN (2018). Knowledge and Practice of Standard Precautions by Health care workers in a Tertiary Health Institution in Enugu, Nigeria. *Nigerian Journal of clinical practice* 21:2 pg149-154.
- [6] Atif, M. L., Brenet ,A., Hageaux .S., Fave M. H., Cochet. C., Baticle, E. (2013). Awareness of standard precautions for 4439 healthcare professionals in 34 institutions in France. *Med Mal Infect*;43:10-6 *Infectious Diseases* 14: e1106–14. *International Journal of Infection Control* 8: 1–7.
- [7] Sreedharan J., Muttappillymyalil J., Venkatramana M. (2011). Knowledge about standard precautions among university hospital nurses in the United Arab Emirates. *East Mediterr Health J* 2011;17:331-4.
- [8] Amoran O, Onwube O.(2013). Infection control and practice of standard precautions among healthcare workers in Northern Nigeria. *J Glob Infect Dis* 2013;5:156-63.
- [9] Parmeggiani C., Abbate R., Marinelli P., Angelillo I. F. (2010). Healthcare workers and health care-associated infections: Knowledge, attitudes, and behavior in emergency departments in Italy. *BMC Infect Dis*; 10:35.
- [10] Punia ,S., Nair S., Shetty, R. S. (2014). Healthcare workers and standard precautions perceptions and determinants of compliance in the emergency and trauma triage of a tertiary care hospital in South India. *International School of Research Notices* 2014:5. [Doi: 10.1155/2014/685072].
- [11] Mukherjee, S, Bhattacharyya, A, SharmaSarkar, B, Goswami, DN, Ghosh,S, Samanta A,(2012). Knowledge and practice of standard precautions and awareness regarding post-exposure prophylaxis for HIV among interns of a medical college in West Bengal, India. *International Journal of Infection Control* 15: 1–7.
- [12] Tobin E. A. Asogun D. A., Odia I., Ehidihamhen G. (2013) Knowledge and practice of infection control among health workers in a tertiary hospital in Edo State, Nigeria. *Direct Research Journal of Health Pharmacy*. 1(2): 20 – 27.
- [13] Abubakar, S. M. Haruna, H., Teryila, K. R., Hamina, D., Ahmadu, I, Babaji, M., (2015). Assessment of knowledge and practice of standard precautions among nurses working at Federal Medical Centre Gombe, Nigeria. *Direct Research Journal of Health Pharmacy*. 3(1): 1-11.
- [14] Adinma, .ED., Ezeama, C., and Asuzu , M.C. (2009). Knowledge and practice of universal precautions against blood borne pathogens amongst house officers and nurses in tertiary health institutions in Southeast Nigeria. *Nigerian Journal of Clinical Practice* 77: 1079–1082..
- [15] Reda ,A.A., Fisseha, S., Mengistie, B, and Vandeweerd, J-M. (2010). Standard precautions: occupational exposure and behavior of health care work-ers in Ethiopia. *PLoS One* 5: e14420.
- [16] Luo ,Y, He, G., Zhou, J., and Luo, Y. (2010). Factors impacting compliance with standard precautions in nursing, China. *International Journal of Infectious Diseases* 14: e1106–14.

The background of the top section is a dark red color with a light red grid pattern. A large, faint, light red caduceus symbol is centered in the background. To the left of the caduceus, there is a faint ECG line. In the bottom right corner, there is another faint ECG line.

AD Publications

Sector-3, MP Colony, Bikaner, Rajasthan, INDIA

www.adpublications.org, www.imjhealth.org, info@imjhealth.org