

International Multispeciality Journal of Health

ISSN: 2395-6291



www.imjhealth.org

Volume-9, Issue-7, July 2023

Preface

We would like to present, with great pleasure, the inaugural volume-9, Issue-7, July 2023, of a scholarly journal, *International Multispecialty Journal of Health*. This journal is part of the AD Publications series *in the field of Medical, Health and Pharmaceutical Research Development*, and is devoted to the gamut of Medical, Health and Pharmaceutical issues, from theoretical aspects to application-dependent studies and the validation of emerging technologies.

This journal was envisioned and founded to represent the growing needs of Medical, Health and Pharmaceutical as an emerging and increasingly vital field, now widely recognized as an integral part of scientific and technical statistics investigations. Its mission is to become a voice of the Medical, Health and Pharmaceutical community, addressing researchers and practitioners in below areas

Clinical Specialty and Super-specialty Medical Science:

It includes articles related to General Medicine, General Surgery, Gynecology & Obstetrics, Pediatrics, Anesthesia, Ophthalmology, Orthopedics, Otorhinolaryngology (ENT), Physical Medicine & Rehabilitation, Dermatology & Venereology, Psychiatry, Radio Diagnosis, Cardiology Medicine, Cardiothoracic Surgery, Neurology Medicine, Neurosurgery, Pediatric Surgery, Plastic Surgery, Gastroenterology, Gastrointestinal Surgery, Pulmonary Medicine, Immunology & Immunogenetics, Transfusion Medicine (Blood Bank), Hematology, Biomedical Engineering, Biophysics, Biostatistics, Biotechnology, Health Administration, Health Planning and Management, Hospital Management, Nephrology, Urology, Endocrinology, Reproductive Biology, Radiotherapy, Oncology and Geriatric Medicine.

Para-clinical Medical Science:

It includes articles related to Pathology, Microbiology, Forensic Medicine and Toxicology, Community Medicine and Pharmacology.

Basic Medical Science:


It includes articles related to Anatomy, Physiology and Biochemistry.

Spiritual Health Science:

It includes articles related to Yoga, Meditation, Pranayam and Chakra-healing.

Each article in this issue provides an example of a concrete industrial application or a case study of the presented methodology to amplify the impact of the contribution. We are very thankful to everybody within

that community who supported the idea of creating a new Research with *IMJ Health*. We are certain that this issue will be followed by many others, reporting new developments in the Medical, Health and Pharmaceutical Research Science field. This issue would not have been possible without the great support of the Reviewer, Editorial Board members and also with our Advisory Board Members, and we would like to express our sincere thanks to all of them. We would also like to express our gratitude to the editorial staff of AD Publications, who supported us at every stage of the project. It is our hope that this fine collection of articles will be a valuable resource for *IMJ Health* readers and will stimulate further research into the vibrant area of Medical, Health and Pharmaceutical Research.



Dr. Kusum Gaur
(Chief Editor)



Mr. Mukesh Arora
(Managing Editor)

Board Members

Dr. Kusum Gaur (Editor-in-chief)

Dr. Kusum Gaur working as professor Community Medicine and member of Research Review Board of Sawai Man Singh Medical College, Jaipur (Raj) India.

She has awarded with WHO Fellowship for IEC at Bangkok. She has done management course from NIHFWS. She has published and present many research paper in India as well as abroad in the field of community medicine and medical education. She has developed Socio-economic Status Scale (Gaur's SES) and Spiritual Health Assessment Scale (SHAS). She is 1st author of a book entitled " Community Medicine: Practical Guide and Logbook.

Research Area: Community Medicine, Biostatistics, Epidemiology, Health and Hospital Management and Spiritual Health.

Mukesh Arora (Managing Editor)

BE (Electronics & Communication), M.Tech (Digital Communication), currently serving as Assistant Professor in the Department of ECE.

Dr. AMER A. TAQA

Dr. AMER A. TAQA is Professor and Head in Dental Basic Science Mosul University, Mosul, IRAQ. He has been registrar of department of Dental Basic Science Mosul University, Mosul, IRAQ. He has published about 100 of research papers and out of that 50 were of international level. He has awarded many times for scientific researches by Government. He has been member of many examination committees and also is a Member in Iraqi Scientific Staff. He has been working as Editor - reviewer in many journals.

Research Area: Dental Science.

Dr. I.D. Gupta

Dr. I. D. Gupta is Professor Psychiatry and working as additional Principal and Dean of student welfare in SMS Medical College, Jaipur.

He is recipient of Prof. Shiv Gautam oration award by Indian Psychiatric Society. He has done training in YMRS at Monte Carlo and BPRS at Singapore. He has been President Indian Psychiatric Society, Rajasthan State Branch. He is author of "Psycho Somatic Disorder" chapter in 1st edition post graduate text book of Psychiatry by Vyas and Ahuja. He has also worked with National Mental Health Programme and has a lot of publication.

Research Area: Community Mental Health, Psycho somatic and liaison Psychiatry.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY

Dr. Anuradha Yadav

Dr. Anuradha Yadav is working as Professor Physiology, SMS Medical College, Jaipur (Rajsthan) India. She is a popular medical teacher and research scholar who had many publications in indexed journals.

Research Area: CVS & CNS physiology, Medical Education and Spiritual Health.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practcal Guide and Logbook".

Research Area: His area of Interest are Epidemiology, Biostatistics and Spiritual Health.

Prof. Dillip Kumar Parida

Professor and Head in the Department of Oncology, AIIMS, Bhubaneswar.

He has done the Professional Training in Japan (Osaka University, NIBI, AHCC Research Association, Hyogo Ion Beam Center), ESTRO Fellowship in Denmark and India(AIIMS Delhi, BARC Mumbai, SCB Medical College-Cuttak, MKCG Medical College-Berhampur).

Research Area: Brachytherapy, Total Skin Electron Irradiation, Palliative Radiotherapy, Stereotactic & Conformal radiotherapy, Radiation Cell Biology, Cancer Genetics.

Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY.

Dr Rajeev Sharma (MS; FMAS; FIMSA;FCLS)

He is working as Professor, Department of Surgery, Government Medical College, Chandigarh, India. He has done FMAS, FIMSA and FCLS along with MS (Gen Surgery).

He has about 50 international and national publications to his credit. He has held various positions in the Association of Minimal Access Surgeons of India (AMASI) from time to time. He has also acted as instructor of various AMASI skill courses held at different places in India. He has established Surgical Technique learning centre at GMCH Chandigarh for imparting training to the budding surgeons in the field of minimal access surgery. He is also the reviewer in the subject in various journals.

Research Area: Minimal Access Surgery.

Dr Anshu Sharma (MS ANATOMY)

She is Presently working as assistant professor in the department of Anatomy, GMCH, Chandigarh. She has many publications in various national and international journals. She is executive member of Anatomical Society of India (ASI) and North Chapter of ASI. She is also a member of editorial board of Journal of Medical College Chandigarh.

Research Area: Congenital Malformation, Developmental Anatomy.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practical Guide and Logbook".

Research Area: His areas of Interest are Epidemiology, Biostatistics and Spiritual Health.

Dr. Dilip Ramlakhyani

Dr. Dilip Ramlakhyani working as Associate professor Pathology and member of IT Committee of Sawai Man Singh Medical College, Jaipur (Raj) India. He has published many articles in indexed journals.

Dr. Virendra Singh

Dr. Virendra Singh worked as Supernatant and head of department of Pulmonary Medicine, SMS Medical College, Jaipur (Rajsthan) India.

He has gone abroad for many training courses and to present research papers. He had been chairman of Research Review Board of SMS Medical College, Jaipur. He is a great research scholar and had published book related to his faculty and had many publications in indexed journals.

Dr. Mahesh Sharma

Dr. Mahesh Sharma is a Principle specialist General Surgery in Rajasthan State Government, India. He has been PMO of district hospitals for more than 15 years. He has gone abroad as observer of many of training related to his speciality. He has published and present many research paper in India as well as abroad.

He has developed Spiritual Health Assessment Scale (SHAS) with Dr. Kusum Gaur.

Research Area: General Surgery, Health and Hospital management and Spiritual Health.

Dr. Ravindra Manohar

Professor Community Medicine, working as head of department of PSM, SMS Medical College, Jaipur (Rajsthan) India.

Previously he has worked in BP Kiorala Institute of Medical Sciences, Nepal. He has visited CDC Atlántica for a Statistical workshop. He has been convener of MBBS and PG exams. He is a research scholar and had many publications in indexed journals.

Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Table of Contents

Volume-9, Issue-7, July 2023

S.No	Title	Page No.
1	<p>The Cognitive Burden of Bipolar Disorder: A Case Report of A Patient Managed Successfully using Endoxifen</p> <p>Authors: Dr Mrinmay Kumar Das</p> <p> DOI: https://dx.doi.org/10.5281/zenodo.8198623</p> <p> Digital Identification Number: IMJH-JUL-2023-3</p>	01-04
2	<p>Implementation of Health Education and Home Visits to Adult Patient's Hypertension</p> <p>Authors: Fransiska Keron Ola, Uswatul Khasanah, Dewi Anggraini</p> <p> DOI: https://dx.doi.org/10.5281/zenodo.8211183</p> <p> Digital Identification Number: IMJH-JUL-2023-4</p>	05-13

The Cognitive Burden of Bipolar Disorder: A Case Report of A Patient Managed Successfully using Endoxifen

Dr Mrinmay Kumar Das

Senior Consultant - Behavioural Medicines, Jaypee Hospital, Sector 128, Noida 201304, Uttar Pradesh, India.

Received:- 25 June 2023/ Revised:- 11 July 2023/ Accepted: 25 July 2023/ Published: 31-07-2023

Copyright © 2023 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— *Bipolar disorder carries a substantial long-term risk, and can lead to cognitive impairment. This can have far-reaching impacts on both social and professional realms of life, thus contributing to increased disability. This case report describes a patient with bipolar disorder who was treated with lithium and suffered worsening cognition. Switching to endoxifen led to an alleviation of symptoms and restoration of cognitive function. It also enabled the discontinuation of lithium therapy within 4 weeks of initiating endoxifen. The case highlights the potential of endoxifen to be used as a primary mood stabilizer in patients with bipolar disorder in the presence of cognitive function.*

Keywords— *Endoxifen, bipolar disorder, lithium, cognitive dysfunction.*

I. INTRODUCTION

Bipolar disorder is known to be a severe and debilitating condition, that affects 2.4% of people worldwide. The long-term psychological risk of bipolar disorder is substantial. Cognitive impairment is known to occur during mood states as well as during euthymia and leads to functional impairment.¹ Furthermore, it has been suggested that cognitive impairment is a marker of bipolar disorder at onset, and is detectable in the early years of bipolar disorder.² Apart from the direct impact of bipolar disorder on cognition, medications used for the management of bipolar disorder can also impact cognition. Lithium is one such mood stabilizer that is known to negatively impact cognition in the short term and in the long term.¹

We present a case of a patient with bipolar disorder on treatment with lithium carbonate, who reported worsening cognition. Switching from lithium carbonate to endoxifen restored cognitive function, making endoxifen a useful option in the management of bipolar disorder.

II. CASE REPORT

A 38-year-old male with known bipolar disorder was on maintenance treatment with lithium 800 mg per day, and thyroxine (for hypothyroidism). He presented with complaints of inability to make business decisions. The patient was self-employed, and involved in running a family-owned packaging business. The patient worked for 2 to 3 days per week, instead of 6 days per week. Along with cognitive dysfunction, the patient then presented with sleeplessness, angry behaviour, and restlessness for two weeks. The recent changes in his behavior were not appreciated by some of the employees. The patient's wife helped with the business and business-related decisions when the patient was unable to cope.

The patient had a first manic episode six years prior, and the patient required injectables and brief hospitalization. At the time, the patient was discharged with lithium carbonate as the primary mood stabilizer. Another major episode occurred two years prior, when he developed hypothyroidism, likely due to treatment with lithium. The patient was subsequently initiated on 25 micrograms of thyroxine, while lithium treatment was continued. Following this, the patient was stable on treatment with lithium carbonate and thyroxine, with only mild depressive symptoms such as irritability and reluctance to participate in social events.

History-taking revealed significant disability in paying attention. The patient reported that he faced difficulty making effective business decisions and in holding meaningful conversations. His family members, including his wife, endorsed these difficulties in social situations, such as the inability to decide on a car parking arrangement for a birthday party. His indecisiveness appeared to be a form of cognitive problems and had become a challenge for his business. Evaluation of symptoms was carried out using the Cognitive Complaints in Bipolar Disorder Rating Assessment (COBRA) scale. This 16-item self-reported instrument measures subjective cognitive dysfunctions including executive function, processing speed, working memory, verbal learning and memory, attention/concentration, and mental tracking. A 4-point scale is used to rate each item (0=never, 1=sometimes, 2=often, and 3=always). The scores of each item are added to give the total score, and higher scores indicate more subjective complaints.³ The assessment of symptoms revealed a score of 24, and the patients had severe functional impairment. The patient was very reluctant to try other medications, including anti-depressants.

The patient was started on endoxifen 8 mg, while lithium was slowly tapered and stopped within 4 weeks. Within 8 weeks, the patient was stable and manic symptoms resolved. He started managing his business better, and there were no complaints of disruptive behavior. After two months of follow-up, the patient seemed calmer and more cooperative, as well as more thoughtful. His ability to participate in therapeutic and social conversations improved. The COBRA score at follow-up was 6. The patient stated that he felt “a lot clearer in the head”.

The patient’s wife reported that he was managing the business better, and social engagement had improved. Furthermore, compared to before endoxifen treatment, the patient did not have more trouble remembering recent occurrences and recalling recent conversations, did not need more assistance with transport, and did not have more difficulty in finding the right words to speak.

III. DISCUSSION

The case described in this report showcases the role of endoxifen in the management of bipolar disorder, based on its favorable pharmacodynamics and safety profile. It was effective in replacing lithium as the main mood stabilizer, as it led to remission, and also improved cognitive function. Bipolar disorder is a leading cause of disability, with both patients and their families experiencing lower functional status, reduced quality of life, and stress on personal relationships, as well as loss of employment, difficulty in regaining employment, and absenteeism from work. These collectively contribute to the cost and disability of the disease.⁴

Cognitive dysfunction is linked to adverse psychosocial outcomes and unemployment outcomes.⁵ Cognitive impairment is not just a characteristic of bipolar disorder but impacts disease outcomes as well. Patients with bipolar disorder experience defects in primary attention processing, executive function, and verbal memory.⁶ It has been reported that 12–40% of patients with bipolar disorder have specific deficits in verbal learning, working memory, and executive function, while a similar proportion have impairments in all cognitive domains.⁷ Studies indicate that impaired verbal memory is linked to the duration of illness, the number of manic episodes, and the number of psychiatric hospitalizations.⁶ Since functional deficits persist beyond the manic stage, it increases the direct and indirect costs associated with bipolar disorder.⁵ Health-related absenteeism is significantly higher among patients with bipolar disorder than those without (18.9 days vs. 7.4 days annually), and it also increases health benefit costs.⁸

Caregiver burden is another aspect of bipolar disease and involves the expenditure of time and money. It is also the cause for worry, tension, and grief. This caregiving strain has been reported for various psychiatric illnesses, affecting over 90% of caregivers of those with bipolar disorder when the patient is hospitalized, and persisting in 70% of caregivers at 15 months after hospitalization.⁹

One aspect of concern is the cognitive dysfunction arising from the use of lithium for the management of bipolar disorder. Lithium has displayed a potential for thyroid, renal and cognitive dysfunction.¹⁰ As seen in this case, the patient developed hypothyroidism after long-term use of lithium and then reported cognitive dysfunction. A meta-analysis has reported that lithium treatment is associated with impairment of immediate verbal learning and memory, as well as creativity.¹¹ Impaired

cognitive function has clinical implications for the treatment of bipolar disorder, as these adverse effects are a leading cause of medication nonadherence.¹²

Endoxifen is a protein kinase C (PKC) inhibitor that has demonstrated anti-manic activity in patients with bipolar disorder I. It effectively reduces the Young Mania Rating Scale (YMRS) score and improves the Montgomery–Åsberg Depression Rating Scale (MADRS) score, leading to early remission from the disease.¹³ The advantage of endoxifen is the safety profile, which is reflected in the low rate of treatment discontinuations due to adverse effects (0-0.9%),^{13,14} in contrast with the high rate of treatment discontinuation with lithium (54% of patients report discontinuing lithium on at least one occasion, of which 62% cite adverse effects as the reason for discontinuation).¹⁵ In addition, endoxifen does not negatively impact cognitive function.^{13,14} The adverse effects of endoxifen are mild to moderate, and resolve within the same day.¹⁴ Long-term use of endoxifen is well-tolerated, and it is a useful agent for preserving functioning and improving quality of life.¹⁶

IV. CONCLUSION

Neurocognitive difficulties in bipolar disorder are known to occur with lithium treatment. In this case, neurocognitive difficulties due to lithium therapy impacted the patient's ability to run his business as well as his profession. Treating the patient with endoxifen led to improvement in the patient's neurocognitive dysfunction, and improvement of manic symptoms. This case also described an additional benefit of endoxifen including increased work productivity and reduced absenteeism, better decision-making, and reduction of caregiver burden. Furthermore, the side effects noted with lithium (hypothyroidism) were not noted with endoxifen. Overall, the benefits of endoxifen indicate that it is an ideal choice for the management of bipolar disorder.

REFERENCES

- [1] Saches M, Bauer IE, Galvez JF, Zunta-Soares GB, Soares JC. (2015). The management of cognitive impairment in bipolar disorder: Current status and perspectives. *American Journal of Therapeutics*. 22(6): 477–486. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4424179/>.
- [2] Misowiak KW, Lewandowski KE. (2020). Trajectory of cognitive functions in bipolar disorder: For better or worse? *Acta Psychiatrica Scandinavica*. 141: 95–97. <https://onlinelibrary.wiley.com/doi/full/10.1111/acps.13144>.
- [3] Rosa AR, Mercade C, Sanchez-Moreno J, Sole B, del Mar Bonnin C, Torrent C, et al. (2013). Validity and reliability of a rating scale on subjective cognitive deficits in bipolar disorder (COBRA). *Journal of Affective Disorders*. 150(1): 29-36. <https://www.sciencedirect.com/science/article/abs/pii/S0165032713001821>.
- [4] Stimmel GL. (2004). Economic grand rounds: the economic burden of bipolar disorder. *Psychiatry Online*. 55(2): 117-118. <https://ps.psychiatryonline.org/doi/10.1176/appi.ps.55.2.117>.
- [5] Sole B, Jimenez E, Torrent C, Reinares M, del Mar Bonnin C, Torres I, et al. (2017). Cognitive impairment in bipolar disorder: Treatment and prevention strategies. *International Journal of Neuropsychopharmacology*. 20(9): 670-680. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5570032/>.
- [6] Vrabie M, Marinescu V, Talasman A, Tautu O, Drima E, Miclutia I. (2015). Cognitive impairment in manic bipolar patients: important, understated, significant aspects. *Annals of General Psychiatry*. 14: 41. <https://annals-general-psychiatry.biomedcentral.com/articles/10.1186/s12991-015-0080-0>.
- [7] Xu N, Huggon B, Saunders KEA. (2019). Cognitive impairment in patients with bipolar disorder: Impact of pharmacological treatment. (2019) *CNS Drugs*. 34: 29-46. <https://link.springer.com/article/10.1007/s40263-019-00688-2>.
- [8] Gardner HH, Kleinman NL, Brook RA, Rajagopalan K, Brizee TJ, Smeeding JE. (2006). The economic impact of bipolar disorder in an employed population from an employer perspective. *Journal of Clinical Psychiatry*. 67(8): 1209-1218. https://www.psychiatrist.com/wp-content/uploads/2021/02/12928_economic-impact-bipolar-disorder-employed-population.pdf.
- [9] Perlick DA, Rosenheck RA, Mcklowitz, DJ, Kaczynski R, Ketter T, Wisniewski W, et al. (2008). Caregiver burden and health in bipolar disorder A cluster analytic approach. *Journal of Nervous and Mental Disease*. 196(6): 484-491. <http://doi.org/10.1097/NMD.0b013e3181773927>.
- [10] Rybakowski JK. (2018). Challenging the negative perception of lithium and optimizing its long-term administration. *Frontiers in Molecular Neuroscience*. 11: 349. <https://www.frontiersin.org/articles/10.3389/fnmol.2018.00349/pdf>.
- [11] Wingo AP, Wingo TS, Harvey PD, Baldessarini RJ. (2009). Effects of lithium on cognitive performance: A meta-analysis. *Journal of Clinical Psychiatry*. 70(11): 1588-1597. <https://www.psychiatrist.com/jcp/bipolar/effects-lithium-cognitive-performance-meta-analysis/>.
- [12] Pachet AK, Wisniewski AM. (2003). The effects of lithium on cognition: An updated review. *Psychopharmacology*. 170: 225-234. <https://link.springer.com/article/10.1007/s00213-003-1592-x>.

- [13] Ahmed A, Sheikh S, Reddy MS, Reddy MS, Prasad BSV, Verma KK, et al. (2016). Endoxifen, a new treatment option for mania: A double-blind, active-controlled trial demonstrates the antimanic efficacy of endoxifen. *Clinical and Translational Science*. 9(5): 252–9. <https://ascpt.onlinelibrary.wiley.com/doi/10.1111/cts.12407>.
- [14] Ahmed A, Sheikh S, Khan MA, Chaturvedi A, Patel P, Patel R, et al. (2021). Endoxifen: A new, protein kinase C inhibitor to treat acute and mixed mania associated with bipolar I disorder. *Bipolar Disorders*. 23(6): 595–603. <https://onlinelibrary.wiley.com/doi/10.1111/bdi.13041>.
- [15] Ohlund L, Ott M, Oja S, Bergqvist M, Lundqvist R, Sandlund M, et al. (2018). Reasons for lithium discontinuation in men and women with bipolar disorder: a retrospective cohort study. *BMC Psychiatry*. 18: 37. <https://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-018-1622-1>.
- [16] Garg S. (2022). First report of endoxifen treatment for 1 year: A case report of bipolar disorder. *International Multispecialty Journal of Health*. 8(5): 1-4. <https://dx.doi.org/10.5281/zenodo.6596685>

Implementation of Health Education and Home Visits to Adult Patient's Hypertension

Fransiska Keron Ola¹, Uswatul Khasanah², Dewi Anggraini³

^{*1}Nursing Department, Dirgahayu College of Health Sciences Samarinda, Jalan Pasundan No. 21, Samarinda, 75122, Kalimantan Timur, Indonesia

^{2,3}Nursing Department, University of Muhammadiyah Jakarta, Jalan Cempaka Putih Tengah I No.1, Kota Jakarta Pusat, 10510, Indonesia

*Corresponding Author

Received: - 07 July 2023/ Revised: - 16 July 2023/ Accepted: 24 July 2023/ Published: 31-07-2023

Copyright © 2023 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract— Hypertension is a public health issue and a silent risk for cardiovascular disease. The goal of this study is of health education sessions and home visits to reduce blood pressure in patients and adults with uncontrolled hypertension. The study is a cluster randomized controlled trial was performed. The trial will be conducted on 40 individuals aged ≥ 36 to 45 years old with hypertensive (with systolic BP ≥ 140 mmHg and diastolic BP ≥ 90 mmHg and patients with uncontrolled blood pressure were equally and randomly allocated into 2 groups. We provide health education sessions with the syllabus of the American Heart Association with modification of booklet and a home visits. The period of intervention is 12 weeks. 2 weeks each month. The participants of the control received only usual care. SPSS 22 programs utilize to analyze the findings, using the analysis of covariance.

Results: The level of knowledge hypertension is 80% with Low level pre interventions and post-intervention with knowledge of 80% High level. This shows that statistically there is a significant effect of Health Education interventions through Home Visits on the level of knowledge (p -value 0.000; <0.05). The difference in the level of knowledge after the intervention with the control group was obtained (p -value 0.00; <0.05). Health Education intervention based on Home Visit to Self Efficacy (p -value 0.000; <0.05).

Conclusion: The results showed that health education and home visits were very effective in increasing patient knowledge about hypertension and reducing blood pressure in patients with hypertension.

Keywords— Implementation, Health Education, Home Visits, Adult Patient's, Hypertension.

I. INTRODUCTION

World Health Organization (WHO) data for 2018 shows that around 1.13 billion people in the world have hypertension, meaning that 1 out of 3 people in the world is diagnosed with hypertension. The number of people with hypertension continues to increase every year, it is estimated that by 2025 there will be 1.5 billion people affected by hypertension, and it is estimated that every year 10.44 million people die from hypertension and its complications.

(1). Community-based health education programs can assist in improving health outcomes in patients with chronic diseases (2). Hypertension is a silent disease of the masses with an increasing prevalence and poor control (3) Empowering nurses to manage hypertension at the community level is feasible with positive good probability results for patients . The worldwide burden of hypertension contributes significant risk of heart failure, coronary artery events, stroke, renal failure, disability, and premature death. Modifiable lifestyle behaviors such as tobacco use, physical inactivity, unhealthy diet, and alcohol abuse are major risk factors contributing to the increased incidence of high blood pressure. A population-based approach to lowering blood pressure levels in the general population even at modest levels has the potential to substantially reduce morbidity and mortality and possibly delay the onset of hypertension (4).

II. MATERIALS AND METHODS

2.1 Study Design

The study used quasi-experimental design with a cluster randomized controlled trial.

2.2 Intervention Protocols

We randomly assigned 2 groups, namely the intervention group and the control group. The intervention group received door-to-door health education while the control group received regular health care. After that in session 1 the nurse made a home visit to the intervention patient to discuss previous basic knowledge, experience, obstacles, difficulties, misunderstandings, behavior, treatment, and lifestyle of people with hypertension, and create a comfortable environment between participants and nurses. In the second session, the nurse measures blood pressure and provides an intervention pretest questionnaire to assess the client's level of knowledge regarding hypertension using the Hypertension Knowledge Level Scale (HK-LS) questionnaire.

3rd Session Nurses visit homes with hypertension and provide health education about hypertension from house to house using booklets modified from the guidelines for the Hypertension Syllabus from the American Heart Association, namely the definition of hypertension, factors that cause hypertension, signs and symptoms of hypertension, complications of hypertension. Management of hypertension care, and treatment of hypertension. Nurses also discuss personal/family problems they face in controlling blood pressure, remind them of physical activity and fruit and vegetable intake, suggest visiting health care facilities if blood pressure is not within the normal range, and provide feedback to families after counseling. After that, the nurse gave a post-test knowledge questionnaire about hypertension using the Hypertension Knowledge Level Scale (HK-LS) questionnaire. The nurse measures blood pressure one week later after the counseling and the nurse measures the patient's behavior after the counseling using a self-efficacy questionnaire.

2.3 Sampling and Sample Size

The implementation of this study was with the population of the Jawa District community in RT 19 and RT 25 with the final adult population aged 36-45 years. The inclusion criteria for this study were aged 36-45 years, could speak Indonesian well, had a history of uncontrolled hypertension of 140/90 mmHg, and were currently on anti-hypertension medication, and available contact numbers such as mobile phones. The exclusion criteria were pregnant women, blind or hearing impaired people, bedridden patients, and participants diagnosed with kidney disease, cancer, heart disease, chronic obstructive pulmonary disease, and mental illness. There were 40 participants, namely 20 people in the control group and 20 people in the intervention group.

2.4 Instruments

The instrument used was a knowledge questionnaire about hypertension from the Hypertension Knowledge Level Scale (HK-LS). As well as the Behavior questionnaire from the self-efficacy questionnaire. The variable measured is the level of knowledge about hypertension using a questionnaire and assessed using an ordinal scale with a range of results (0-21) low knowledge, (22-43) moderate knowledge, and (44-66) high knowledge. Self-efficacy questionnaires were assessed using an ordinal scale with a range of (0-6) obedient behavior, (7-10) moderately obedient behavior, and (11-14) obedient behavior. Blood pressure measurement using a digital sphygmomanometer measuring instrument brand Omron with a range of 120-130/80-85 Normal and > 180/110 Severe hypertension.

2.5 Ethical Consideration

This study was approved by Jawa District Primary Health care Pasundan and Jawa District No. 03/Kel. Jawa/011/2022 for voluntary participants, informed consent, and confidentiality of participant's identity.

2.6 Data collection and Data analysis

The author uses primary data sources from the residents of Jawa District with predetermined criteria. Participants fill out the approval form to follow the research implementation process.

Analysis data used the Statistical Program for Social Science (SPSS) version 22 which included *Wilcoxon*, and *Mann-Whitney*. Difference Test Analysis.

III. RESULTS AND DISCUSSION

3.1 Results

Data was collected by visiting participants who had hypertension. The age of participants was 36 to 45 years old. The total of intervention participants is 20 and the control participants are 20.

The majority age 45 years of respondents 12 people (30%). Female 26 people (65%). The education of respondent is high school 22 people (55%). The majority of respondents had been diagnosed with hypertension for 1-5 years as many as 24 people (60%).

3.1.1 Age Variable

The frequency distribution of the age of the respondents in the intervention group and the group can be seen in table 1 as follows:

TABLE 1
AGE FREQUENCY DISTRIBUTION OF AGGREGATE ADULT RESPONDENTS AGE 36-45 YEARS OLD.

Characteristics	Intervention		Control		Total	
	n	%	n	%	n	%
36 Years Old	3	15%	4	20%	7	17,5%
37 Years Old	1	5%	0	0%	1	2,5%
38 Years Old	0	0%	1	5%	1	2,5%
39 Years Old	0	0%	1	5%	1	2,5%
40 years Old	3	15%	1	5%	4	10%
41 Years Old	0	0%	3	15%	3	7,5%
42 years Old	2	10%	2	10%	4	10%
43 Years Old	4	20%	0	0%	4	10%
44 Years Old	1	5%	2	10%	3	7,5%
45 years Old	6	30%	6	30%	12	30%
Total	20	100	20	100	40	100

(Source: Primary data for 2022)

The result of the frequency distribution of the age of the respondents in the intervention group and the group can be seen in table 1 with Frequency Distribution of Aggregate Adult Respondents Age 36-45 years old with 45 Years old (30%) 6 people intervention and variable control 6 people (30%). (Table 1)

3.1.2 Gender Variable

The frequency distribution of the gender of the respondents in the intervention group and the group can be seen in table 2 as follows:

TABLE 2
DISTRIBUTION OF AGGREGATE GENDER FREQUENCY OF RESPONDENTS AGGREGATE ADULTS AGE 36-45 YEARS OLD.

Characteristics	Intervention		Control		Total	
	n	%	n	%	n	%
Man	2	10%	12	60%	14	35%
Woman	18	90%	8	40%	26	65%
Total	20	100	20	100	40	100

(Source: Primary data for 2022)

The result of gender distribution of the respondents can be seen it shows that the majority of respondents are female, amounting to 26 people (65%). (Table 2)

3.1.3 Education Level Variable

The frequency distribution of respondents' education level in the intervention and control groups can be seen in table 3 as follows:

TABLE 3
FREQUENCY DISTRIBUTION OF AGGREGATE AGGREGATE EDUCATION LEVELS OF ADULT RESPONDENTS
EDUCATION 36-45 YEARS OLD.

Characteristics	Intervention		Control		Total	
	n	%	n	%	n	%
Junior high school	6	30%	6	30%	12	30%
Senior High School	11	55%	11	55%	22	55%
Bachelor degree	3	15%	3	15%	6	15%
Total	20	100	20	100	40	100

(Source: Primary data for 2022)

The result of education level of distribution of respondents' education can be seen in it shows that the majority of respondents have a secondary education level, totaling 22 people (55%). The majority of respondents in this study had basic education (elementary school) as many as 10 people (44%). While the second respondent has high school education as many as 6 people (30%). (Table 3).

3.1.4 Variable Diagnosed with Hypertension

The distribution of length of time diagnosed with hypertension in the intervention and control groups can be seen in table 4 as follows:

TABLE 4
FREQUENCY DISTRIBUTION OF AGE OLD DIAGNOSED WITH HYPERTENSION AGGREGATE RESPONDENTS
AGGREGATE ADULTS AGE 36-45 YEARS OLD.

Characteristics	Intervention		Control		Total	
	n	%	n	%	n	%
1-5 Years	12	60%	12	60%	24	60%
6-10 Years	7	35%	7	35%	14	35%
>11 Years	1	5%	1	5%	2	5%
Total	20	100	20	100	40	100

(Source: Primary data for 2022)

The result of variable diagnosed with hypertension distribution of length of time diagnosed with hypertension to respondents can be seen it shows that the majority of respondents have been diagnosed with hypertension for 1-5 years as many as 24 people (60%). (Table 4)

3.1.5 Level of Knowledge about Hypertension

The distribution of hypertension knowledge in the intervention and control groups can be seen in table 5 as follows:

TABLE 5
DISTRIBUTION OF EMPLOYMENT FREQUENCY OF RESPONDENTS AGGREGATE ADULTS AGED 36-45 YEARS OLD.

Characteristics	Intervention		Control		Total	
	n	%	n	%	n	%
Housewife	16	80%	7	35%	23	57,5%
Seamstress	1	5%	1	5%	2	5%
Teachers	1	5%	2	15%	3	7,5%
Private	2	10%	10	50%	12	30%
Total	20	100	20	100	40	100

(Source: Primary data for 2022)

The results of the knowledge showed that differences in the level of knowledge about hypertension of respondents before and after the Home Visit-based Health Education intervention showed that the majority of respondents had a high level of knowledge of 16 respondents (80%), but there were respondents who had a moderate level of knowledge of 5 respondents (20%). This shows that statistically there is a significant effect of Health Education interventions through Home Visits on the level of knowledge (p-value 0.000; <0.05). (Table 5)

3.1.6 Level of Blood Pressure

The distribution of blood pressure in the intervention group can be seen in table 6 as follows:

TABLE 6
DISTRIBUTION OF AGGREGATE BLOOD PRESSURE OF ADULT RESPONDENTS AGE 36-45 YEARS OLD.

Category	Blood pressure				Category	Blood pressure			
	Pre systole		Post systole			pre diastole		Post diastole	
	n	%	n	%		n	%	n	%
130	0	0%	7	35%	80	0	0%	9	45%
140	6	30%	6	30%	90	10	50%	10	50%
150	6	30%	6	30%	100	9	45%	1	5%
160	5	25%	0	0%	110	1	5%		
170	0	0%	1	5%					
180	2	10%	0	0%					
190	1	5%	0	0%					
Total	20	100	20	100	Total	20	100	20	100

(Source: Primary data for 2022)

The distribution of blood pressure in respondents showing that the majority of respondents had pre-test systolic blood pressure of 140-190 mmHg while post-test systolic blood pressure decreased to 130-150 mmHg. The pre-test diastolic blood pressure ranged from 90-110 mmHg while the post-test diastolic blood pressure was 80-90 mmHg (Table 6).

3.1.7 Level of Anthropometry Abdominal Circumference

Anthropometric distribution of abdominal circumference in the intervention and control groups can be seen in table 7 as follows:

TABLE 7
ANTHROPOMETRIC DISTRIBUTION (ABDOMINAL CIRCUMFERENCE) IN AGGREGATE ADULT RESPONDENTS AGED 36-45 YEARS OLD.

Category	Abdominal Circumference				Category	Abdominal Circumference			
	Pre intervention		Post intervention			Control pre		Control Post	
	n	%	n	%		n	%	n	%
Not obese (< 80 cm)	6	30%	7	35%	Not obese (< 80 cm)	6	30%	0	0%
obesity (>80 cm)	14	70%	13	65%	obesity (>80 cm)	14	70%	14	70%
Total	20	100	20	100	Total	20	100	20	100

(Source: Primary data for 2022)

The result of the Body Mass Index respondents showed that the majority of respondents were obese, 16 people (80%) in the intervention group and 14 people (70%) in the control group (Table 7).

3.1.8 Level of Body Mass Index

The distribution of Body Mass Index in the intervention and control groups can be seen in table 8 as follows:

TABLE 8
DISTRIBUTION OF BODY MASS INDEX IN AGGREGATE ADULT RESPONDENTS AGED 36-45 YEARS OLD.

Category	Body Mass Index				Category	Body Mass Index			
	Pre intervention		Post intervention			Control Pre		Control Post	
	n	%	n	%		n	%	n	%
Underweight (BB< 18,5 kg/m ²)	0	0%	0	0%	Underweight BB< 18,5 kg/m ²)	1	5%	1	5%
Normal (BB 18,5-24,9 kg/m ²)	3	15%	2	10%	Normal (BB 18,5-24,9 kg/m ²)	3	15%	3	15%
Excess body (BB > 25 kg/m ²)	1	5%	2	10%	Excess bodies (BB > 25 kg/m ²)	14	70%	14	70%
Obesity (BB > 30 kg/m ²)	16	80%	16	80%	Obesity (BB > 30 kg/m ²)	2	10%	2	10%
Total	20	100	20	100	Total	20	100	20	100

(Source: Primary data for 2022)

The result of distribution Body Mass Index to respondents can be seen in showing that the majority of respondents were obese, 16 people (80%) in the intervention group and 14 people (70%) in the control group. (Table 8)

3.1.9 Level of Knowledge Levels Hypertension

The distribution of Knowledge Levels about Hypertension in the intervention and control groups can be seen in table 9 as follows:

TABLE 9
DISTRIBUTION OF KNOWLEDGE LEVELS ABOUT HYPERTENSION IN AGGREGATE RESPONDENTS OF ADULTS AGE 36-45 YEARS OLD.

Characteristics	Knowledge level			
	Pre		Post	
	n	%	n	%
Rendah (0-21)	22	55%	16	40%
Sedang (22-43)	18	45%	8	20%
High Level of Knowledge (44-66)	0	0%	16	40%
Total	40	100	40	100

(Source: Primary data for 2022)

The result distribution of the knowledge level of respondents in the intervention and control groups can be seen in table 9. It shows that the majority of respondents had a low level of knowledge before the intervention, totaling 22 people (55%) and experienced an increase after the intervention, totaling 16 people (40%) had a high level of knowledge. (Table 9)

3.1.10 Level of Differences in Knowledge before and after the intervention

The distribution of Knowledge Levels about Hypertension in the intervention group before and after the action can be seen in table 10 as follows:

TABLE 10
DISTRIBUTION OF KNOWLEDGE LEVELS ABOUT HYPERTENSION BEFORE AND AFTER INTERVENTION IN AGGREGATE RESPONDENTS OF ADULTS AGE 36-45 YEARS OLD.

Characteristic	Knowledge level				Value
	Pre		Post		<i>p-value</i>
	n	%	n	%	
Low knowledge (0-21)	6	30%	0	0%	0,000
Currently (22-43)	14	70%	4	20%	
high knowledge (44-66)	0	0%	16	80%	
Total	20	100	20	100	

(Source: Primary data for 2022)

The results of differences in the level of knowledge about hypertension of respondents before and after the Home Visit-based Health Education intervention can be seen in table 10 showing that the majority of respondents have a high level of knowledge as many as 16 respondents (80%), but there are respondents who have a moderate level of knowledge as many as 4 respondents (20%). This shows that statistically there is a significant effect of Health Education interventions through Home Visits on the level of knowledge (pvalue 0.000; <0.05). (Table 10).

IV. DISCUSSION

The existence of genetic factors in certain families will cause that family to have a risk of suffering from hypertension. This is associated with an increase in intracellular sodium levels and a lower ratio of potassium to sodium. Someone who has parents with hypertension is twice as likely to suffer from hypertension than someone who does not have a family history of hypertension. In addition, 70-80% of essential hypertension cases are found with a family history of hypertension. Based on the theory above, the researchers synthesized that there were genetic factors that led to an increased incidence of hypertension cases in adults. From the data above, it can be concluded that a very significant increase in cases of hypertension has occurred almost all over the world, especially cases of hypertension occurring in the late adult age group. The cause of the increased

incidence of hypertension cases is due to genetics. Hypertension tends to increase, especially in those aged over 40 years (5). The older a person is, the regulation of calcium metabolism becomes disrupted, resulting in large amounts of calcium circulating in the blood vessels. Blood will become denser and blood pressure increases, calcium deposits in the walls of blood vessels cause narrowing so that blood flow is disrupted and triggers an increase in blood pressure (6).

There are also large studies as well assess the relationship between measurements BMI with disease incidence associated with cardiovascular disease (hypertension), indicating that waist circumference measurement and/or waist-to-hip ratio have a better degree of accuracy in estimating the degree of obesity and the risk of cardiovascular disease. Where is abdominal obesity (central) and visceral obesity have an important role in the pathogenesis occurrence of hypertension. Research others have also shown that compared to the population in Europe, Asian populations have a tendency to gain fat abdominal viscera and insulin resistance which is higher at BMI levels that do not show fibrous tissue such as in hypertensive patients (7). Nurses performing relevant interventions are included teaching/guidance/counseling in lifestyle modification changes, medications and procedures such as timing and dosage as well as drug and physical activity interactions, and cases management. After the home visit, follow-up was done via telephone call performed every two weeks by a trained nurse. During follow-up, Trained nurses monitor previous health problems and current patient conditions, as well as modifications in them knowledge, behavior, and status (3).

Health education using the syllabus from health education from the American Hearth Association are modified by using a Hypertension buckle let accompanied by home visits and controlling blood pressure checks, Showed that health education and home visits were very effective in increasing patient knowledge about hypertension and lowering blood pressure in patients with uncontrolled hypertension in adults.

V. CONCLUSIONS

The level of knowledge hypertension is 80% with Low level pre interventions and post-intervention with knowledge of 80% High level. This shows that statistically there is a significant effect of Health Education interventions through Home Visits on the level of knowledge (p-value 0.000; <0.05). The difference in the level of knowledge after the intervention with the control group was obtained (p-value 0.00; <0.05). Health Education intervention based on Home Visit to Self Efficacy (p-value 0.000; <0.05).

VI. RECOMMENDATIONS

Based on the implementation of community nursing practice that has been carried out, there are several suggestions addressed to:

6.1 Development of nursing knowledge

Nursing science education can develop special nursing care guidelines for hypertension in adult aggregates in families and communities using integrated nursing theory. The guide is expected to involve the participation of families and communities to support changes in adult behavior, especially aggregate late adults so that the activities designed can be more comprehensive in efforts to prevent complications of hypertension.

6.2 For the Health Service

Providing a special program for controlling hypertension in late-adult aggregates aged 36-45 years and providing facilities and infrastructure for prevention and control of hypertension in late-adult aggregates.

6.3 Community Health Care Center (Puskesmas)

Providing health promotion media about hypertension with audio and visual for hypertensive patients in adult aggregates and dissemination through social media such as What Sapp and other media to special Community Health Workers (Kader) of hypertension in adult aggregates aged 36-45 years.


ACKNOWLEDGEMENT

We would like to thank especially the Head of the Nursing Department in Sekolah Tinggi Ilmu Kesehatan Dirgahayu Samarinda and to Nursing Faculty at Universitas Muhammadiyah Jakarta, Indonesia.

REFERENCES

[1] K. T. Mills *et al.*, "Comparative effectiveness of implementation strategies for blood pressure control in hypertensive patients: A

- systematic review and meta-analysis," *Ann. Intern. Med.*, vol. 168, no. 2, pp. 110–120, 2018.
- [2] T. H. Jafar *et al.*, "Multicomponent intervention versus usual care for management of hypertension in rural Bangladesh, Pakistan and Sri Lanka: Study protocol for a cluster randomized controlled trial," *Trials*, vol. 18, no. 1, pp. 1–12, 2017.
- [3] N. Srinivasapura Venkateshmurthy *et al.* 2018. Power Heart Project - a Nurse Care Coordinator Led, mHealth Enabled Intervention to Improve the M," *Trials*. (19) 1 : 1–9.
- [4] C. H. Lu *et al.*, "Community-based interventions in hypertensive patients: A comparison of three health education strategies," *BMC Public Health*, vol. 15, no. 1, pp. 1–9, 2015.
- [5] M. K. Khanal *et al.*, "Effectiveness of community-based health education and home support program to reduce blood pressure among patients with uncontrolled hypertension in Nepal: A cluster-randomized trial," *PLoS One*, vol. 16, no. 10 October, pp. 1–17, 2021.
- [6] G. Ogedegbe *et al.*, "HHS Public Access," vol. 129, no. 20, pp. 2044–2051, 2015.
- [7] D. Neupane, C. S. McLachlan, B. Christensen, A. Karki, H. B. Perry, and P. Kallestrup, "Community-based intervention for blood pressure reduction in Nepal (COBIN trial): Study protocol for a cluster-randomized controlled trial," *Trials*, vol. 17, no. 1, pp. 1–7, 2016.
- [8] N. Widjaya, F. Anwar, R. Laura, R. R. Puspawati, and E. Wijayanti, "Hubungan Usia Dengan Kejadian Hipertensi di Kecamatan Kresek dan Tegal Angus , Kabupaten Tangerang The Association Between Age and Incidences of Hypertention in Kresek District And Tegal Angus District , Tangerang Regency," vol. 26, no. 3, pp. 131–138, 2018.
- [9] S. Fatmawati, J. Jafriati, and K. Ibrahim, "Hubungan Life Style Dengan Kejadian Hipertensi Pada Usia Dewasa (20-44 Tahun) Di Wilayah Kerja Puskesmas Puuwatu Kota Kendari Tahun 2017," *J. Ilm. Mhs. Kesehat. Masy. Unsyiah*, vol. 2, no. 6, p. 198151, 2017.
- [10] J. J. Gonidjaya, B. J. Que, N. E. Kailola, C. R. Titaley, and I. Kusadhiani, "Prevalensi Dan Karakteristik Penderita Hipertensi Pada Penduduk Desa Banda Baru Kabupaten Maluku Tengah Tahun 2020 Central Maluku District 2020," vol. 3, no. April, 2021.



AD Publications

Sector-3, MP Colony, Bikaner, Rajasthan, INDIA

www.adpublications.org, www.imjhealth.org, info@imjhealth.org