

International Multispeciality Journal of Health

ISSN: 2395-6291



www.imjhealth.org

Volume-9, Issue-6, June 2023

Preface

We would like to present, with great pleasure, the inaugural volume-9, Issue-6, June 2023, of a scholarly journal, *International Multispecialty Journal of Health*. This journal is part of the AD Publications series *in the field of Medical, Health and Pharmaceutical Research Development*, and is devoted to the gamut of Medical, Health and Pharmaceutical issues, from theoretical aspects to application-dependent studies and the validation of emerging technologies.

This journal was envisioned and founded to represent the growing needs of Medical, Health and Pharmaceutical as an emerging and increasingly vital field, now widely recognized as an integral part of scientific and technical statistics investigations. Its mission is to become a voice of the Medical, Health and Pharmaceutical community, addressing researchers and practitioners in below areas

Clinical Specialty and Super-specialty Medical Science:

It includes articles related to General Medicine, General Surgery, Gynecology & Obstetrics, Pediatrics, Anesthesia, Ophthalmology, Orthopedics, Otorhinolaryngology (ENT), Physical Medicine & Rehabilitation, Dermatology & Venereology, Psychiatry, Radio Diagnosis, Cardiology Medicine, Cardiothoracic Surgery, Neurology Medicine, Neurosurgery, Pediatric Surgery, Plastic Surgery, Gastroenterology, Gastrointestinal Surgery, Pulmonary Medicine, Immunology & Immunogenetics, Transfusion Medicine (Blood Bank), Hematology, Biomedical Engineering, Biophysics, Biostatistics, Biotechnology, Health Administration, Health Planning and Management, Hospital Management, Nephrology, Urology, Endocrinology, Reproductive Biology, Radiotherapy, Oncology and Geriatric Medicine.

Para-clinical Medical Science:

It includes articles related to Pathology, Microbiology, Forensic Medicine and Toxicology, Community Medicine and Pharmacology.

Basic Medical Science:

It includes articles related to Anatomy, Physiology and Biochemistry.

Spiritual Health Science:

It includes articles related to Yoga, Meditation, Pranayam and Chakra-healing.

Each article in this issue provides an example of a concrete industrial application or a case study of the presented methodology to amplify the impact of the contribution. We are very thankful to everybody within

that community who supported the idea of creating a new Research with *IMJ Health*. We are certain that this issue will be followed by many others, reporting new developments in the Medical, Health and Pharmaceutical Research Science field. This issue would not have been possible without the great support of the Reviewer, Editorial Board members and also with our Advisory Board Members, and we would like to express our sincere thanks to all of them. We would also like to express our gratitude to the editorial staff of AD Publications, who supported us at every stage of the project. It is our hope that this fine collection of articles will be a valuable resource for *IMJ Health* readers and will stimulate further research into the vibrant area of Medical, Health and Pharmaceutical Research.



Dr. Kusum Gaur
(Chief Editor)



Mr. Mukesh Arora
(Managing Editor)

Board Members

Dr. Kusum Gaur (Editor-in-chief)

Dr. Kusum Gaur working as professor Community Medicine and member of Research Review Board of Sawai Man Singh Medical College, Jaipur (Raj) India.

She has awarded with WHO Fellowship for IEC at Bangkok. She has done management course from NIHFWS. She has published and present many research paper in India as well as abroad in the field of community medicine and medical education. She has developed Socio-economic Status Scale (Gaur's SES) and Spiritual Health Assessment Scale (SHAS). She is 1st author of a book entitled " Community Medicine: Practical Guide and Logbook.

Research Area: Community Medicine, Biostatistics, Epidemiology, Health and Hospital Management and Spiritual Health.

Mukesh Arora (Managing Editor)

BE (Electronics & Communication), M.Tech (Digital Communication), currently serving as Assistant Professor in the Department of ECE.

Dr. AMER A. TAQA

Dr. AMER A. TAQA is Professor and Head in Dental Basic Science Mosul University, Mosul, IRAQ. He has been registrar of department of Dental Basic Science Mosul University, Mosul, IRAQ. He has published about 100 of research papers and out of that 50 were of international level. He has awarded many times for scientific researches by Government. He has been member of many examination committees and also is a Member in Iraqi Scientific Staff. He has been working as Editor - reviewer in many journals.

Research Area: Dental Science.

Dr. I.D. Gupta

Dr. I. D. Gupta is Professor Psychiatry and working as additional Principal and Dean of student welfare in SMS Medical College, Jaipur.

He is recipient of Prof. Shiv Gautam oration award by Indian Psychiatric Society. He has done training in YMRS at Monte Carlo and BPRS at Singapore. He has been President Indian Psychiatric Society, Rajasthan State Branch. He is author of "Psycho Somatic Disorder" chapter in 1st edition post graduate text book of Psychiatry by Vyas and Ahuja. He has also worked with National Mental Health Programme and has a lot of publication.

Research Area: Community Mental Health, Psycho somatic and liaison Psychiatry.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY

Dr. Anuradha Yadav

Dr. Anuradha Yadav is working as Professor Physiology, SMS Medical College, Jaipur (Rajsthan) India. She is a popular medical teacher and research scholar who had many publications in indexed journals.

Research Area: CVS & CNS physiology, Medical Education and Spiritual Health.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practcal Guide and Logbook".

Research Area: His area of Interest are Epidemiology, Biostatistics and Spiritual Health.

Prof. Dillip Kumar Parida

Professor and Head in the Department of Oncology, AIIMS, Bhubaneswar.

He has done the Professional Training in Japan (Osaka University, NIBI, AHCC Research Association, Hyogo Ion Beam Center), ESTRO Fellowship in Denmark and India(AIIMS Delhi, BARC Mumbai, SCB Medical College-Cuttak, MKCG Medical College-Berhampur).

Research Area: Brachytherapy, Total Skin Electron Irradiation, Palliative Radiotherapy, Stereotactic & Conformal radiotherapy, Radiation Cell Biology, Cancer Genetics.

Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Dr. Lokendra Sharma

Dr. Lokendra Sharma is Associate Professor Pharmacology and working as Nodal officer of SMS Medical College, Jaipur.

He is recipient of WHO Fellowship award on Poison Patient Management at Vietnam. He is resource faculty for Experimental Toxicology and Basic Course for Medical Education. He is presented and published a lot of research articles at national and international level.

Research Area: PHARMACOLOGY.

Dr Rajeev Sharma (MS; FMAS; FIMSA;FCLS)

He is working as Professor, Department of Surgery, Government Medical College, Chandigarh, India. He has done FMAS, FIMSA and FCLS along with MS (Gen Surgery).

He has about 50 international and national publications to his credit. He has held various positions in the Association of Minimal Access Surgeons of India (AMASI) from time to time. He has also acted as instructor of various AMASI skill courses held at different places in India. He has established Surgical Technique learning centre at GMCH Chandigarh for imparting training to the budding surgeons in the field of minimal access surgery. He is also the reviewer in the subject in various journals.

Research Area: Minimal Access Surgery.

Dr Anshu Sharma (MS ANATOMY)

She is Presently working as assistant professor in the department of Anatomy, GMCH, Chandigarh. She has many publications in various national and international journals. She is executive member of Anatomical Society of India (ASI) and North Chapter of ASI. She is also a member of editorial board of Journal of Medical College Chandigarh.

Research Area: Congenital Malformation, Developmental Anatomy.

Dr. Rajeev Yadav

Dr. Rajeev Yadav is working as Associate Professor Community Medicine, SMS Medical College, Jaipur (Rajsthan) India. He is member of Research Review Board of the Institute.

He has authored a book entitled "Community Medicine: Practical Guide and Logbook".

Research Area: His areas of Interest are Epidemiology, Biostatistics and Spiritual Health.

Dr. Dilip Ramlakhyani

Dr. Dilip Ramlakhyani working as Associate professor Pathology and member of IT Committee of Sawai Man Singh Medical College, Jaipur (Raj) India. He has published many articles in indexed journals.

Dr. Virendra Singh

Dr. Virendra Singh worked as Supernatant and head of department of Pulmonary Medicine, SMS Medical College, Jaipur (Rajsthan) India.

He has gone abroad for many training courses and to present research papers. He had been chairman of Research Review Board of SMS Medical College, Jaipur. He is a great research scholar and had published book related to his faculty and had many publications in indexed journals.

Dr. Mahesh Sharma

Dr. Mahesh Sharma is a Principle specialist General Surgery in Rajasthan State Government, India. He has been PMO of district hospitals for more than 15 years. He has gone abroad as observer of many of training related to his speciality. He has published and present many research paper in India as well as abroad.

He has developed Spiritual Health Assessment Scale (SHAS) with Dr. Kusum Gaur.

Research Area: General Surgery, Health and Hospital management and Spiritual Health.

Dr. Ravindra Manohar

Professor Community Medicine, working as head of department of PSM, SMS Medical College, Jaipur (Rajsthan) India.

Previously he has worked in BP Kiorala Institute of Medical Sciences, Nepal. He has visited CDC Atlántica for a Statistical workshop. He has been convener of MBBS and PG exams. He is a research scholar and had many publications in indexed journals.



Dr. Praveen Mathur

Dr. Praveen Mathur is working as Professor- Pediatric Surgery and is recipient of Commonwealth Fellowship in Pediatric Laparoscopy from Uk and fellowship award in minimal access Surgery (FMAS). He has done Clinical observer ship in the Department of Pediatric Surgery, Johns Hopkins University, Baltimore, USA. (2008). He has presented and published a number of research articles at national and international level. He is reviewer of prestigious Journal of Pediatric Surgery (JPS) and World Journal of Gastroenterology, Journal of neonatal Surgery (JNS).

Research Area: Pediatric Surgery & Laparoscopy.

Table of Contents

Volume-9, Issue-6, June 2023

S.No	Title	Page No.
1	<p>Prescribing Trends in Bipolar I Disorder and Usage of Endoxifen: An Indian Perspectivez</p> <p>Authors: M S Reddy, Sameer Malhotra, Aswin Ajit, Jyoti Kapoor, U Gauthamadas, Anoop Vincent, Meena Gnanasekharan, Satyakant Trivedi, Pawan Adatia, Arnab Ghosh Hazra, N Rangarajan, Vishal Sawant</p> <p> DOI: https://dx.doi.org/10.5281/zenodo.8098731</p> <p> Digital Identification Number: IMJH-JUN-2023-4</p>	01-03

Prescribing Trends in Bipolar I Disorder and Usage of Endoxifen: An Indian Perspective

M S Reddy^{1*}, Sameer Malhotra², Aswin Ajit³, Jyoti Kapoor⁴, U Gauthamadas⁵, Anoop Vincent⁶, Meena Gnanasekharan⁷, Satyakant Trivedi⁸, Pawan Adatia⁹, Arnab Ghosh Hazra¹⁰, N Rangarajan¹¹, Vishal Sawant¹²

¹MD, DPM, Asha Hospital, Hyderabad, Andhra Pradesh, India, 500034

²MD, DM, DNB, Max Hospital, Saket, Delhi, India, 110017

³MD, DPM, Dhanya Doctor's Chamber, Ernakulam, Kerala, India, 682032

⁴MBBS, DPM, Paras Hospital, Sushant Lok, Gurgaon, Haryana, India, 122002

⁵MBBS, MD, Doc Gautham's Neuro Center—"We re-mind your brain", Chennai, Tamil Nadu, India, 600031

⁶MD, DM, Sree Narayana Medical College, Cochin, Kerala, India, 683594

⁷MD, DM, Reach Psychiatry Centre, Bangalore, Karnataka, India, 560043

⁸MD, DM, Bansal Hospital, Bhopal, Madhya Pradesh, India, 462016

⁹MBBS, DPM, Aasha Hospital Ayurvedic Layout, Nagpur, Maharashtra, India, 440008

¹⁰MD, Zenith Super Speciality Hospital, Kolkata, West Bengal, India, 700056

¹¹MD, DPM, Psymed Hospital, Chennai, Tamil Nadu, India, 600031

¹²MD, DPM, Parooques Poly Clinic, Mumbai, Maharashtra, India, 400092

*Corresponding Author

Received:- 23 May 2023/ Revised:- 07 June 2023/ Accepted: 18 June 2023/ Published: 30-06-2023

Copyright © 2023 International Multispecialty Journal of Health

This is an Open-Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted Non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Abstract—

Objective: Bipolar I disorder (BD-I) is a psychiatric illness characterized by erratic moods and impulsive behaviors. This survey assesses the prescription preferences among Indian psychiatrists for BD-I, current unmet needs, and benefits of endoxifen (a direct protein kinase C inhibitor) as a mood stabilizing agent in BD-I patients.

Methods: A literature review was carried out based on data from the PubMed Database to identify relevant articles (published between January 1980 and May 2022) using specific keywords. Twenty clinically relevant questions belonging to six major domains were drafted: (i) key attributes in the selection of psychotropic medications as front-line therapy; (ii) preference for mania and mixed episodes in front-line therapy; (iii) key attributes for the selection of maintenance therapy; (iv) preference for maintenance therapy; (v) role and positioning of endoxifen in front-line and maintenance settings; and (vi) determination of patient subgroups who can experience benefits from endoxifen. A total of 77 psychiatrists with significant experience in managing patients with bipolar disorder were identified across different cities in India. An electronic survey link to the questionnaire was sent to all the participants to record their views.

Results: For BD-I mania, a combination of mood stabilizers and an atypical antipsychotic was preferred. Typical side effects noted in Indian BD-I patients on lithium, valproate, or carbamazepine therapy include drug-induced tremors, hepatic failure, and metabolic disturbances. Experts suggested endoxifen in patients with acute and severe BD-I mania due to its good efficacy and tolerability profile. For the management of mixed episodes of BD-I, experts preferred endoxifen in combination with an antipsychotic therapy or selective serotonin reuptake inhibitor. Maintenance therapy was suggested in patients with more relapse tendencies and after a severe manic episode that warrants hospitalization.

Conclusion: The good tolerability profile of endoxifen encourages its use in patients whose current treatment options for BD-I bring challenging side effects.

Keywords— Bipolar I disorder, prescribing practices, management, endoxifen, India.

I. INTRODUCTION

Bipolar disorder (BD) is a chronic recurrent psychiatric illness associated with significant disability and high suicide risk [1]. It is characterized by mood instability (mania, depression, or both) and impulsive behaviors [2-4]. Bipolar disorder encompasses the following subtypes [5,6]:

- Bipolar I disorder (BD-I) is defined by manic episodes that last at least 1 week (most of the day, nearly every day) or by severe manic symptoms that require immediate hospitalization. Manic episodes may be preceded by and may be followed by hypomanic or major depressive episodes (**Figure 1**).
- Bipolar II disorder (BD-II) is defined by a pattern of hypomanic episodes lasting at least 4 consecutive days (most of the day, nearly every day) and major depressive episodes typically lasting at least 2 weeks.
- Cyclothymic disorder is defined by recurrent hypomanic and depressive episodes for 2 years or longer.




FIGURE 1: Symptoms of mania and mania with mixed features

Adapted from: DSM-5 (Fifth edition) [5], National Institute of Mental Health: Bipolar Disorder [6], and Hu J et al. 2014 [7].

Bipolar disorder is common in clinical psychiatric practice. A large-scale cross-sectional survey across 11 countries in the USA, Europe, and Asia (World Mental Health Initiative) indicated that the lifetime prevalence of bipolar spectrum disorders was 2.4%, with a prevalence of 0.6% for BD-I and 0.4% for BD-II [8]. In India, the prevalence of bipolar spectrum disorder is 0.1%, of which BD-I accounts for roughly 20% of the patient population [9]. Studies from India suggest a substantially higher incidence of mania than major depressive episodes in the course of BD-I [9,10]. Drugs typically used for the treatment of BD-I mania include mood stabilizers (MS) and antipsychotics as monotherapy or a combination of both in patients with severe illness and those requiring hospitalization [11]. For the management of BD-I depression, initial treatment options include antipsychotics, MS, and antidepressants [11]. A systematic review by Vázquez GH highlighted a disease recurrence risk in BD patients (96.0% BD-I; mean onset age: 23.1 years; follow-up: 1.9 years) during treatment with an MS or antipsychotic drug as 39.3% (21.9%/year) [12]. Combination therapy is more likely than monotherapy to result in treatment discontinuation because of problems with tolerability, extrapyramidal symptoms, sedation, and weight gain [13]. Since the last decade, studies have recognized the major role of overactive protein kinase C (PKC) intracellular signaling in psychiatric disorders such as BD and schizophrenia [14-19]. Endoxifen (4-OH-N-desmethyltamoxifen), a direct PKC inhibitor, was found to be safe and effective

in managing acute mania, severe mania with psychotic features, and mixed episodes of BD-I [20-23]. In this article, we aim to capture prescribing preferences among Indian psychiatrists for BD-I and current unmet needs in the medical management of BD-I. We also aim to gather insights from experts on their experience and benefits of endoxifen as a mood-stabilizing agent in patients with BD-I.

II. METHODOLOGY

2.1 Literature Review and Development of the Questionnaire

An extensive literature review was carried out based on data from the PubMed Database to identify relevant articles published between January 1980 and May 2022 using keywords such as “bipolar,” “bipolar 1 disorder,” “mania,” “hypomania,” “depression,” “mixed state,” “mood stabilizers,” “antipsychotics,” “anticonvulsants,” “antidepressants,” “protein kinase C,” “mixed episodes,” “treatment,” “maintenance,” “endoxifen,” “guidelines,” and “management.” Various combinations of keywords were used. Twenty clinically relevant questions were drafted (**Figure 2**). The content of the questionnaire was prepared keeping in mind the practice environment in India (**Supplementary file**).




FIGURE 2: Major domains of the survey questionnaire

BD-I: Bipolar I disorder.

2.2 Study Design

For the phase I survey, psychiatrists (N=77) with at least 10 years of clinical experience in managing BD patients across 56 locations in India were identified. Around 64 psychiatrists started endoxifen therapy in BD-I patients at various time points in a 2-year study duration (2021–2022). The questionnaire was rolled out to the participants through an online survey platform. One-to-one online discussions were held with each of the participants and preferences were captured. Respondents were expected to fill in their preferences for a given indication from the options provided. Certain open-ended questions were included in the survey to seek experts' opinions based on clinical practice.

2.3 Data Analysis

Descriptive statistics using the frequency counts of the responses to each question were generated. For data analysis, no formal statistical modeling was done. Here, we report interim results of the phase I survey (n=77). The survey will be extended further (phases II and III; 2023 onward) to include opinions from psychiatrists across different cities in India.

III. RESULTS

3.1 Clinician's Perspective on the Current Burden of BD-I in India

Based on the clinical practice, experts agreed that around 5%–29% of their patient population are diagnosed with BD (84%; N=64 experts). More than 50% of BD patients are diagnosed to have BD-I (45%; N=34 experts).

3.2 Current Challenges in the Medical Management of BD-I and the Role of Endoxifen in Front-Line Settings

3.2.1 BD-I Mania

Experts opined that the selection of psychotropic medications against mania in patients with BD-I is dependent on efficacy (faster onset of action and reduced rate of relapse) and safety profile (minimal metabolic disturbances, no weight gain, and no change in blood glucose or thyroid hormone levels). Consideration of age and gender and minimal need for drug monitoring were rated as the least preferred criteria for medication selection. For the treatment of BD-I mania, a combination of an MS (lithium, valproate, or carbamazepine) and an atypical antipsychotic was the preferred first choice for the majority of experts (67%; N=51 experts), followed by an MS (lithium, valproate, or carbamazepine) with a typical antipsychotic (33%; N=25), whereas valproate with benzodiazepine was the third preference (30%; N=23 experts) (**Table 1**). Lithium, endoxifen, olanzapine, endoxifen with atypical antipsychotics, and valproate were the preferred choice of therapy in women of reproductive age (15–49 years). In women of the nonreproductive age group, endoxifen, lithium, and valproate were suggested. Nonadherence to the treatment regimen is more frequent with combination therapy due to tolerability issues and pill burden than with monotherapy. Typical side effects noted in Indian BD-I patients on MS therapy (lithium, valproate, or carbamazepine) include drug-induced tremors, hepatic failure, and metabolic disturbances (**Table 2**). Adherence to the treatment is affected due to sedation and weight gain.

TABLE 1
PREFERENCE FOR MANAGEMENT OF BD-I MANIA (N=77)

	First preference (%)	Second preference (%)	Third preference (%)
Lithium monotherapy	11	9	11
Antipsychotic monotherapy	3	9	13
Divalproex/valproate monotherapy	7	14	12
Lithium with benzodiazepine	1	12	18
Valproate with benzodiazepine	5	9	30
Carbamazepine/oxcarbamazepine with and without benzodiazepine	1	1	8
MS (lithium, valproate, carbamazepine/oxcarbamazepine) in combination with atypical antipsychotic	67	12	3
MS (lithium, valproate, carbamazepine/oxcarbamazepine) in combination with typical antipsychotic	5	33	5

MS: Mood stabilizer.

TABLE 2
LIST OF SIDE EFFECTS AND ADVERSE EVENTS EXPERIENCED WITH THE USAGE OF LITHIUM, CARBAMAZEPINE, AND DIVALPROEX/VALPROATE IN INDIAN PATIENTS WITH BD-I MANIA

	Lithium	Divalproex/valproate	Carbamazepine
Side effects	<ul style="list-style-type: none"> Tremor Nausea Diarrhea Weight gain Acne Impaired memory Hypothyroidism Delirium Excessive urination Increased serum creatinine levels Metallic taste 	<ul style="list-style-type: none"> Alopecia Weight gain Tremor Hepatotoxicity Gastrointestinal disturbances Liver dysfunction Menstrual irregularities Hyperammonemia Hyponatremia 	<ul style="list-style-type: none"> Agranulocytosis Ataxia Skin rash Weight gain Nausea Withdrawal seizures Elevations in liver enzymes Hepatotoxicity
Caution	Therapeutic drug monitoring required	<ul style="list-style-type: none"> Monitoring liver function and ammonia levels required Therapeutic drug monitoring required 	Therapeutic drug monitoring required

Experts mentioned that mania is associated with an overactive PKC intracellular signaling, and direct PKC inhibition with endoxifen can help in faster remission of symptoms with no incidence of treatment-emergent depression, tremors, weight gain, or metabolic disturbances. There was no need for therapeutic drug monitoring during endoxifen therapy, unlike lithium, valproate, and carbamazepine therapy. **Table 3** lists key attributes of endoxifen that stood out to experts during the phase I survey. For acute manic episodes, experts preferred endoxifen (**Table 4**) in combination with an MS (66%; N=50 experts) or atypical antipsychotic (62%; N=47 experts). Few respondents preferred to use endoxifen alone in the treatment of mild-to-moderate BD-I mania (37%; N=28 experts).

TABLE 3
KEY ATTRIBUTES OF ENDOXIFEN THAT STOOD OUT TO EXPERTS (N=77)

	Response (%)
Fast response	82
No need for therapeutic drug monitoring unlike lithium and divalproex	79
No adverse effects such as a decrease in platelet count, hair loss, and tremors	79
Convenience (one pill a day)	78
Lesser/no impact on metabolic parameters, such as thyroid hormone levels, blood glucose levels, and weight	78
Reduction in pill burden	74
No incidence of treatment-emergent depression	70

TABLE 4
PREFERENCE FOR THE USE OF ENDOXIFEN IN BD-I MANIA (N=77)

	Preference (%)
In combination with an MS	66
In combination with atypical antipsychotics	62
As monotherapy in mild-to-moderate mania	37
In combination with an MS and atypical antipsychotic	34
In combination with a benzodiazepine	28
In combination with typical antipsychotics	12

BD-I: Bipolar I disorder; MS: Mood stabilizer.

3.2.2 Mixed Episodes of BD-I

Around 57% of experts (N=43) mentioned that they were “very concerned” about the incidence of treatment-emergent mania due to antidepressant therapy in patients with BD-I and would not prescribe it. Some experts (37%; N=28) were “somewhat concerned” about the incidence of treatment-emergent mania due to antidepressant therapy in BD-I and would prescribe it if the benefits outweighed the risk. For the treatment of BD-I patients with acute depression, a combination of MS and antipsychotic was the first preferred choice (**Table 5**) for the majority (43%; N=33 experts), followed by an MS and a selective serotonin reuptake inhibitor (SSRI; 34%; N=26 experts), whereas MS in combination with a tricyclic antidepressant (TCA) was the third preference (28%; N=21 experts). For the management of mixed episodes of BD-I, experts preferred endoxifen in combination with an antipsychotic therapy (68%; N=52 experts) or SSRIs (32%; N=24 experts). Few respondents preferred to use endoxifen alone (41%; N=31 experts) or in combination with a TCA (4%; N=3 experts) for the treatment of mixed episodes of BD-I.

TABLE 5
PREFERENCE FOR MANAGEMENT OF BD-I PATIENTS WITH ACUTE DEPRESSION (N=77)

	First preference (%)	Second preference (%)	Third preference (%)
MS in combination with an SSRI	37	34	26
MS in combination with a TCA	–	16	28
MS in combination with an antipsychotic	43	28	22
MS monotherapy	20	22	24

BD-I: Bipolar I disorder; MS: Mood stabilizer; SSRI: Selective serotonin reuptake inhibitor; TCA: Tricyclic antidepressant.

3.2.3 Determination of Patient Pool That Can Reap Maximum Benefits from Endoxifen Therapy in Front-Line Setting

The experts opined that a good tolerability profile of endoxifen encourages its use in patients (**Table 6**) where current treatment options for BD-I bring challenging side effects, such as lithium toxicity, sedation, and tremors (75%; N=57 experts). Other patient subgroups in whom endoxifen has a promising role include: (i) female patients with BD-I (71%; N=54 experts); (ii) working-class BD-I patients as endoxifen has no sedative effect (71%; N=54 experts); and (iii) patients seeking for reduced pill burden (67%; N=51 experts).

TABLE 6
PATIENT PROFILES WHO CAN BENEFIT FROM ENDOXIFEN THERAPY IN FRONT-LINE SETTINGS (N=77)

	Preference (%)
Patients facing challenges with other treatment options	75
Female patients with BD-I	71
Working-class BD-I patients	71
Patients seeking reduced pill burden	67
Patients with mild-to-moderate mania	61
Patients with mixed episode	51
Patients with severe mania, who are frequently hospitalized	50
Elderly patients with BD-I	47

BD-I: Bipolar I disorder; MS: Mood stabilizer.

3.2.4 Duration of Endoxifen Therapy in Front-Line Settings

In the phase I survey, the maximum duration of endoxifen therapy studied in patients with BD-I varied from 1 month to 12 months (**Figure 3**). This is because the experts (N=64) did not initiate endoxifen treatment in the same month.




FIGURE 3: Maximum duration of endoxifen therapy studied in patients with BD-I (N=64)
BD-I: Bipolar I disorder

3.3 Current Challenges in the Medical Management of BD-I in Maintenance Settings and the Role of Endoxifen

Experts agreed that the selection of psychotropic medications for maintenance in patients with BD-I is strongly dependent on the efficacy and safety profile (minimal metabolic disturbances and thyroid hormone levels). Experts indicated maintenance therapy in the following patient types of BD-I:

- After the first episode of acute BD-I mania (63%; N=48 experts)
- After the first mixed episode of BD-I (70%; N=53 experts)
- In patients with bipolar depression (63%; N=48 experts)
- After subsequent episodes (71%; N=54 experts)
- After any episode severe enough to warrant hospitalization (82%; N=62)
- In patients with more relapse tendency (86%; N=65)

Lithium (34%; N=26 experts) monotherapy was the preferred treatment option in the maintenance settings, followed by valproate (29%; N=22 experts) and carbamazepine/oxcarbazepine (32%; 24 experts) monotherapy (Table 7). The most frequently reported reasons for treatment discontinuation during maintenance therapy in Indian patients with BD-I were metabolic disturbances (weight gain and change in glucose and cholesterol levels), sedation, and risk of extrapyramidal symptoms. Experts (100%; N=77) mentioned that they were interested to try endoxifen in maintenance settings of BD-I. As per the Canadian Network for Mood and Anxiety Treatments (CANMAT) guidelines, medications that have been found to be effective in the acute phase should be continued during the maintenance phase [24]. Thus, endoxifen has the potential to be an effective therapeutic for BD for long-term use. With such an encouraging experience, the experts unanimously agreed to try endoxifen in the maintenance settings of BD-I. The key attributes of endoxifen that make it a preferred choice of therapy under maintenance settings are an encouraging real-world experience (good efficacy and safety profile), convenient dosing, and no therapeutic drug monitoring (86%; N=65 experts).

TABLE 7
PREFERENCE FOR MAINTENANCE TREATMENT OF BD-I (N=77)

	First Preference (%)	Second preference (%)	Third preference (%)
Lithium monotherapy	34	16	13
Valproate monotherapy	25	29	13
Carbamazepine/oxcarbazepine monotherapy	–	13	32
Combination of lithium and an atypical antipsychotic	8	17	14
Combination of valproate and an atypical antipsychotic	21	17	14
Combination of lithium/valproate with an extended-release atypical antipsychotic	12	8	13

BD-I: Bipolar I disorder

IV. DISCUSSION

The main findings of this survey of prescribing practices of Indian psychiatrists for the treatment of BD-I are as follows:

- A combination of an MS and atypical antipsychotic is the preferred treatment option for BD-I mania.
- For the management of BD-I patients with acute depression, experts preferred MS in combination with antipsychotic therapy or an SSRI. Experts were concerned about the incidence of treatment-emergent mania due to antidepressant therapy in patients with BD-I.
- Maintenance therapy was suggested in patients with more relapse tendencies and after a severe manic episode that warrants hospitalization.

The current mainstays for the management of BD-I mania are lithium and divalproex/valproate, which are indirect inhibitors of PKC [25]. Endoxifen is a direct PKC inhibitor with proven efficacy independent of CYP2D6-mediated metabolism in patients with BD-I [20]. Direct PKC inhibition with endoxifen can help in the faster remission of symptoms and reduce the chances of inpatient hospitalizations [22]. Experts suggested endoxifen in patients with acute and severe BD-I mania due to

the good efficacy and tolerability profile of endoxifen. Endoxifen was preferred in patients with mixed features of BD-I in combination with an antipsychotic therapy or an SSRI. **Figure 4** lists the BD-I patient pool who can reap maximum benefits from endoxifen therapy in front-line settings (mania and mixed episodes).

Patients who need faster remission of symptoms	Patients seeking for reduced pill burden	Patients facing challenges with other treatment options
Patients with mild-to-moderate mania	Working-class Patients as endoxifen has no sedative effect	Female Patients with BD-I
Patients with severe mania, who are frequently hospitalized	Patients with mixed episode of BD-I	Elderly Patients with BD-I

FIGURE 4: Role and positioning of endoxifen in the medical management of BD-I in front-line settings
BD-I: Bipolar I disorder

Strengths: The psychiatrists were selected to best represent the breadth of knowledge and clinical expertise in the field from all over India. There was no selection bias.

Limitation: The patient's voice was not included in the consensus process. Endoxifen is a comparatively new drug and its usage and duration captured are still limited as compared to standard-of-care treatment for BD-I.

V. CONCLUSION

BD-I is a psychiatric illness characterized by unstable moods, impulsive behaviors, and reduced quality of life. The selection of psychotropic medications against BD-I is strongly dependent on drug efficacy (faster onset of action and reduced relapse rate) and safety profile (minimal metabolic disturbances, no weight gain, and no change in blood glucose or thyroid hormone levels). Treatment with endoxifen is efficacious in controlling mania and mixed episodes of BD-I, with no adverse effects. The major usage of endoxifen was in combination, except in mild-to-moderate mania. Key attributes of endoxifen that stand out to experts were faster remission of symptoms, reduced pill burden, and no incidences of drug-induced tremors, weight gain, and metabolic disturbances, unlike lithium and valproate therapy. The good tolerability profile of endoxifen encourages its use in a wide spectrum of BD-I patients. As per CANMAT guidelines, medications that have been found to be effective in the acute phase should be continued during the maintenance phase. Thus, endoxifen has the potential to be an effective therapeutic for BD-I for long-term use. With such an encouraging experience, experts unanimously agreed to try endoxifen in the maintenance settings of BD-I.

ACKNOWLEDGEMENT

Dr. M S Reddy as lead author contributed during the course of development of the questionnaire used in this survey and reviewed the outcomes of this consensus. Psychiatrists who participated in the survey mentioned in this publication are Dr. Ashwin Mohan, Dr. Kailash Jhalani, Dr. Manish Jain, Dr. Gaurav Trivedi, Dr. Pratheesh P J, Dr. R K Srivastava, Dr. Fabian Almeida, Dr. Prashant Agarwal, Dr. Rohan Jahagirdar, Dr. Narendra Gupta, Dr. Ajay Dogra, Dr. K P Sharma, Dr. Bir Singh Yadav, Dr. Shijo John Joseph, Dr. Pawan Adatia, Dr. Anay Kshirsagar, Dr. S S Kundu, Dr. RamSharan, Dr. Rahul Chandok, Dr. Pavithra Mony, Dr. Satyakant Trivedi, Dr. Meena Gnanasekharan, Dr. Gautam Saha, Dr. Deepak Mansharmani, Dr. Ashraf Ali, Dr. Tarun Nigam, Dr. Arabinda Brahma, Dr. Jayant Dhake, Dr. Osman Ali, Dr. Bhaskar Prasad, Dr. Niel Shah Dhule, Dr. Phani Bhushan, Dr. Himanshu Sareen, Dr. Amar Shinde, Dr. Manmeet Singh, Dr. Anoop Vincent, Dr. M S Karthik, Dr. U Gauthamadas, Dr. Rohit Garg, Dr. Pramod Gupta, Dr. Rajiv Sharma, Dr. Ashim Chatterjee, Dr. Pankaj Mittal, Dr. A Ramachandran, Dr. Anant Kumar Verma, Dr. K S Jyothi, Dr. N Rangarajan, Dr. Sameer Malhotra, Dr. V V Seshamma, Dr. Pawan Rathi, Dr. Cijo Alex, Dr. T C Ramesh Kumar, Dr. Gautam Amin, Dr. Ravindra Tambe, Dr. K Selvaraj, Dr. Bharat Sarode, Dr. Swapnil Deshmukh, Dr. BSV Prasad, Dr. Vijay Chinchole, Dr. Ankur Singhal, Dr. Joice Geo, Dr. Jyoti Kapoor, Dr. Rajeev Trehan, Dr. Vishal Sawant, Dr. Nitin Dalaya, Dr. Anil Kumar, Dr. Bhagwat Rajput, Dr. Suresh Ninan, Dr. Mahalaxmi, Dr. Karthik Duraisamy, Dr. Vimal Somaiya, Dr. Arnab Ghosh Hazra, Dr. Aswin Krishnan Ajit, Dr. Naazneen Ladak, Dr. Ravi Rana, Dr. Shiv Prasad, and Dr. M S Reddy.

AUTHOR CONTRIBUTIONS

All the authors have contributed to Concepts, Design, definition of intellectual content, data acquisition, data analysis, statistical analysis, and manuscript preparation. Dr. MS Reddy has also contributed to manuscript editing and review.

CONFLICT OF INTEREST

None

FUNDING DETAILS

None

REFERENCES

- [1] E. Vieta, M. Berk, T. G. Schulze, A. F. Carvalho, T. Suppes, J. R. Calabrese, et al, "Bipolar disorders," *Nat Rev Dis Primers*, vol. 4, pp. 18008, Mar 2018.
- [2] G. Marzani, A. Price Neff, "Bipolar disorders: Evaluation and treatment," *Am Fam Physician*, vol. 103, issue 4, pp. 227-239, Feb 2021.
- [3] S. Gautam, A. Jain, M. Gautam, A. Gautam, T. Jagawat, "Clinical practice guidelines for bipolar affective disorder (BPAD) in children and adolescents," *Indian J Psychiatry*, vol. 61, Suppl 2, pp. 294-305, Jan 2019.
- [4] Kattimani S, Subramanian K, Sarkar S, Rajkumar RP, Balasubramanian S, "History of lifetime suicide attempt in bipolar I disorder: Its correlates and effect on illness course," *Int J Psychiatry Clin Pract*, vol. 21, issue 2, pp. 118-24 Jun 2017.
- [5] *Diagnostic and Statistical Manual of Mental Disorders (DSM-5), Fifth edition*. Available https://media.mycme.com/documents/168/dsm-5_bipolar_and_related_diso_41789.pdf. Accessed on: 15 December 2022.
- [6] National Institute of Mental Health. *Bipolar Disorder*. (2023, Feb) Available <https://www.nimh.nih.gov/health/topics/bipolar-disorder>. Accessed on: 17 December 2022.
- [7] J. Hu, R. Mansur, R. S. McIntyre, "Mixed specifier for bipolar mania and depression: Highlights of DSM-5 changes and implications for diagnosis and treatment in primary care," *Prim Care Companion CNS Disord*, vol. 16, issue 2, PCC.13r01599, 2014.
- [8] K. R. Merikangas, R. Jin, J. P. He, R. C. Kessler, S. Lee, N. A. Sampson, et al, "Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative," *Arch Gen Psychiatry*, vol. 68, issue 3, pp. 241-251, Mar 2011.
- [9] S. Karthick, S. Kattimani, R. P. Rajkumar, B. Bharadwaj, S. Sarkar, "Long term course of bipolar I disorder in India: Using retrospective life chart method," *J Affect Disord*, vol. 173, pp. 255-260, Mar 2015.
- [10] S. B. Rangappa, S. Munivenkatappa, J. C. Narayanaswamy, S. Jain, Y. C. J. Reddy "Predominant mania course in Indian patients with bipolar I disorder," *Asian J Psychiatr*, vol. 22, pp. 22-27, Aug 2016.
- [11] R. Jain, A. M. Kong, P. Gillard, A. Harrington, "Treatment patterns among patients with bipolar disorder in the United States: A retrospective claims database analysis," *Adv Ther*, vol. 39, issue 6, pp. 2578-2598, 2022.
- [12] G. H. Vázquez, JN Holtzman, Lolich, Ketter TA, Baldessarini RJ. Recurrence rates in bipolar disorder: Systematic comparison of long-term prospective, naturalistic studies versus randomized controlled trials. *Eur Neuropsychopharmacol*. 2015;25(10):1501-12.
- [13] P. A. Geoffroy, B. Etain, C. Henry, F. Bellivier. "Combination therapy for manic phases: A critical review of a common practice," *CNS Neurosci Ther*, vol.18, issue 12, pp. 957-964, Dec 2012.
- [14] A. Saxena, G. Scaini, D.V. Bavaresco, C. Leite, S. S. Valvassori, A. F. Carvalho, et al, "Role of protein kinase C in bipolar disorder: A review of the current literature," *Mol Neuropsychiatry*, vol. 3, pp. 108-124, Nov 2017.
- [15] H.K. Manji, R. H. Lenox, "Long-term action of lithium: A role for transcriptional and posttranscriptional factors regulated by protein kinase C," *Synapse* vol. 16, pp. 11-28, Jan 1994.
- [16] A. Talaie, M. Pourgholami, H. Khatibi-Moghadam, F. Faridhosseini, F. Farhoudi, A. Askari-Noghani, et al, "Tamoxifen: A protein kinase C inhibitor to treat mania: A systematic review and meta-analysis of randomized, placebo-controlled trials," *J Clin Psychopharmacol*, vol. 3, issue 3, pp. 272-275, Jun 2016.
- [17] L. A. Catapano, H. K. Manji, "Kinases as drug targets in the treatment of bipolar disorder," *Drug Discov Today*, vol. 13, issue 7-8, pp. 295-302, Apr 2008.
- [18] L. S. Carroll, N. M. Williams, V. Moskvina, E. Russell, N. Norton, H. J. Williams, et al, "Evidence for rare and common genetic risk variants for schizophrenia at protein kinase C, alpha," *Mol Psychiatry*, vol. 15, issue 11, pp. 1101-1111, Nov 2010.
- [19] H. Singh, "Endoxifen response in schizoaffective disorder: A case series," *IMJ Health*, vol. 8, issue 8, vol 2, pp. 1-5, Feb 2022.
- [20] A. Ahmad, S. Sheikh, M. A. Khan, A. Chaturvedi, P. Patel, R. Patel, et al, "Endoxifen: A new, protein kinase C inhibitor to treat acute and mixed mania associated with bipolar I disorder," *Bipolar Disord*, vol. 23, issue 6, pp. 595-603, Sep 2021.
- [21] A. Ahmad, S. Sheikh, T. Shah, M. S. Reddy, Bsv. Prasad, K. K. Verma, et al, "Endoxifen, a new treatment option for mania: A double-blind, active-controlled trial demonstrates the antimanic efficacy of endoxifen [published correction appears in *Clin Transl Sci*. 2017;10 (2):117]," *Clin Transl Sci*, vol. 9, issue 5, 252-259, Oct 2016.
- [22] M. Gowda, K. Dewani, "Cost-effectiveness of endoxifen therapy in severe psychotic bipolar I disorder: A case report," *Int J Appl Res*, vol. 8, issue 8, pp. 9-13, 2022.
- [23] V. Thanvi, "Real-world long-term experience on endoxifen in bipolar disorder with psychotic symptoms," *Case Rep Psychiatry*, Vol.2022, pp. 3684181, Jul 2022.
- [24] L. N. Yatham, S. H. Kennedy, S. V. Parikh, A. Schaffer, D. J. Bond, B. N. Frey, et al, "Canadian Network for Mood and Anxiety Treatments (CANMAT) and International Society for Bipolar Disorders (ISBD) 2018 guidelines for the management of patients with bipolar disorder," *Bipolar Disord*, vol. 20, issue 2, pp. 97-170, Mar 2018.
- [25] C. A. Zarate, H. K. Manji, "Protein kinase C inhibitors: Rationale for use and potential in the treatment of bipolar disorder," *CNS Drugs*, vol. 23, issue 7, pp. 569-582, 2009.

SUPPLEMENTARY MATERIAL

PRESCRIBING TRENDS IN BIPOLAR I DISORDER AND USAGE OF ENDOXIFEN: AN INDIAN PERSPECTIVE

Major domains	Questions
Clinician's perspective on the current burden of BD-I in India	<p>1. In your clinical practice, what percentage of patient population are diagnosed with BD?</p> <p>a) Less than 5%</p> <p>b) 5%–14%</p> <p>c) 15%–29%</p> <p>d) 30%–49%</p> <p>e) 50%–69%</p> <p>f) More than 70%</p> <p>2. In your clinical practice, what percentage of BD patients are diagnosed with BD-I?</p> <p>a) Less than 5%</p> <p>b) 5%–9%</p> <p>c) 10%–29%</p> <p>d) 30%–49%</p> <p>e) 50%–69%</p> <p>f) More than 70%</p> <p>g) Not sure</p>
Key attributes in selection of psychotropic medications against mania in patients with BD-I	<p>3. In your expert opinion, what are the key attributes in selection of psychotropic medications against mania in patients with BD-I? (Rank in order: 1—most preferred and 6—least preferred)</p> <p>a) Predominantly efficacy</p> <p>b) Avoidance of treatment-emergent mania and depression</p> <p>c) Both efficacy and safety (minimal metabolic disturbances- no weight gain, no change in blood glucose or thyroid hormone levels)</p> <p>d) Minimal need for drug monitoring</p> <p>e) Faster onset of action and reduced rate of relapse</p> <p>f) Age and gender</p>
<p>Prescribing trends among psychiatrists for BD-I mania</p> <p>Note: Endoxifen is not mentioned in the following options</p> <p>There is a separate section which captures the role and benefits of endoxifen in BD-I mania.</p>	<p>4. What is your preference in management of BD-I mania? (Rank in order of efficacy: 1— most preferred and 8—least preferred)</p> <p>a) Lithium monotherapy</p> <p>b) Antipsychotics monotherapy</p> <p>c) Divalproex/valproate monotherapy</p> <p>d) Lithium with benzodiazepine</p> <p>e) Valproate with benzodiazepine</p> <p>f) Carbamazepine/oxcarbamazepine with and without benzodiazepine</p> <p>g) Mood stabilizer (lithium, valproate, carbamazepine/oxcarbamazepine) in combination with atypical antipsychotic</p> <p>h) Mood stabilizer (lithium, valproate, carbamazepine/oxcarbamazepine) in combination with typical antipsychotic</p> <p>5. Open-ended: What is your preferred treatment choice for BD-I mania in female patients?</p> <p>a) In reproductive age group</p> <p>b) In non-reproductive age group</p>
Current challenges in medical management of BD-I mania	<p>6. Open-ended: What are the most frequent adverse events experienced, with usage of below drugs in BD-I mania?</p> <p>a) Lithium</p> <p>b) Divalproex/valproate</p> <p>c) Carbamazepine</p> <p>d) Atypical antipsychotic (risperidone, olanzapine)</p>
Role and positioning of endoxifen in BD-I mania	<p>7. Mania is associated with overactive protein kinase C intracellular signaling. Would you consider direct PKC inhibitor as a treatment option for mania management in your clinical practice?</p> <p>a) Yes</p> <p>b) No</p>

	<p>8. In your expert opinion, which attributes of the endoxifen treatment stand out to you? (Please select all that apply)</p> <ol style="list-style-type: none"> Only direct PKC inhibitor Fast response Convenience (one pill a day) Reduces pill burden No incidence of treatment-emergent depression No need for therapeutic drug monitoring unlike lithium and divalproex No adverse effects such as decrease in platelet count, hair loss, and tremor Lesser/No impact on metabolic parameters such as thyroid hormone levels, blood glucose levels and weight <p>9. Endoxifen at a daily dose of 8 mg was found to be efficacious and safe in BD-I patients with acute manic episodes as compared to 1000 mg divalproex. Would you prefer to prescribe endoxifen as monotherapy or in combination with other therapies? (Please select all that apply)</p> <ol style="list-style-type: none"> As monotherapy in mild to moderate mania In combination with benzodiazepine In combination with atypical antipsychotics In combination with typical antipsychotics In combination with mood stabilizer In combination with mood stabilizer and atypical antipsychotic
<p>Current challenges in medical management of BD-I depression</p>	<p>10. Are you concerned about incidence of treatment-emergent mania due to antidepressant therapy in patients with BD-I disorder?</p> <ol style="list-style-type: none"> Very concerned, I would not prescribe it Somewhat concerned, I would prescribe if benefits outweighed the risk Not really concerned as it is a rare event Unsure
<p>Prescribing trends among psychiatrists for mixed episodes of BD-I</p> <p>Note: Endoxifen is not in the following options. There is a separate section on role and benefits of endoxifen in management of mixed episodes of BD-I</p>	<p>11. Based on your clinical practice, how would you rate the clinical applicability of the following treatments in BD-I patients and acute depression? (Rank in order of efficacy: 1—most preferred and 4—least preferred)</p> <ol style="list-style-type: none"> Mood stabilizer and a selective serotonin reuptake inhibitor Mood stabilizer and tricyclic antidepressant Mood stabilizer in combination with an antipsychotic therapy Mood stabilizer monotherapy
<p>Role and positioning of endoxifen in mixed episodes of BD-I</p>	<p>12. Based on the available efficacy and safety data of endoxifen (in phase III clinical trial, endoxifen significantly improved YMRS and MADRS scores in treatment of acute mania with or without mixed features in BD-I), how would you recommend endoxifen therapy for the management of mixed episodes of BD-I? (Please select all that applies)</p> <ol style="list-style-type: none"> Endoxifen monotherapy Endoxifen in combination with a selective serotonin reuptake inhibitor (SSRI) Endoxifen in combination with a tricyclic antidepressant Endoxifen in combination with an antipsychotic therapy
<p>Determination of patient pool that can reap maximum benefits from endoxifen therapy in front-line</p>	<p>13. In your opinion, which patient pool would benefit maximum from endoxifen therapy? (Please select all that apply)</p> <ol style="list-style-type: none"> Patients who need faster response Patients seeking for reduced pill burden Patients with mild-to-moderate mania Patients with severe mania, who are frequently hospitalized Patients facing challenges with other treatment options Female patients with BD-I Elderly patients with BD-I Patients with mixed episode Working-class patients as endoxifen has no sedative effect

Duration of endoxifen therapy in front-line	14. Open-ended: In your clinical practice, what is the maximum duration of endoxifen therapy you have tried in patients with BD-I?
Key attributes in selection of maintenance therapy for BD-I	15. In your expert opinion, what are the key attributes for your selection of maintenance therapy for BD-I? (Rank in order: 1—most preferred and 7—least preferred) a) Predominantly safety b) Predominantly efficacy c) Avoidance of treatment-emergent mania and depression d) Both efficacy and safety (minimal metabolic changes, thyroid hormone levels) e) Minimal need for drug monitoring f) Faster onset of action and reduced rate of relapse g) Age and gender
Prescribing trends among psychiatrists for maintenance therapy in BD-I Note: Endoxifen is not in the following options.	16. In which of the following patient types would you suggest/recommend maintenance therapy in BD-I? (Please select all that apply) a) After the first episode of acute mania b) After subsequent episodes c) After the first mixed episode of BD-I d) After any episode severe enough to warrant hospitalization e) In patients with more relapse tendency f) In patients with bipolar depression 17. How would you rate the clinical applicability of the following maintenance therapies in patients with BD-I? (Rank in order of efficacy: 1—most preferred and 6—least preferred) a) Lithium monotherapy b) Valproate monotherapy c) Carbamazepine/oxcarbazepine monotherapy d) Combination of lithium and an atypical antipsychotic e) Combination of valproate and an atypical antipsychotic f) Combination of lithium/valproate with an extended-release atypical antipsychotic g) None of the above
Current challenges in medical management of BD-I in maintenance settings	18. In your clinical practice, what are the most frequently reported reasons for discontinuation during maintenance treatment of BD-I? (Rank in order: 1—most reported and 6— least reported) a) Pill burden b) Alopecia, tremors c) Risk of extrapyramidal symptoms d) Metabolic disturbances (weight gain, change in glucose and cholesterol levels) e) Drug titrations and drug monitoring f) Sedation
Role and benefits of endoxifen in maintenance settings of BD-I	19. In real world settings, endoxifen has > 20,000 Rx in Indian setting (Ref: IMS ORG MAT Feb`22) and the duration of endoxifen treatment is also on increasing trend due to good safety and efficacy profile. With such encouraging experience, would you try using endoxifen in maintenance setting of BD-I? a) Yes b) No 20. According to you, which of the following attributes of endoxifen can make it a preferred choice for maintenance in BD-I? a) Good safety profile b) Convenient dosing and no therapeutic drug monitoring required c) Encouraging real world experience d) Efficacy observed so far in trials may be expected over longer durations of treatment e) All the above

The background of the page is a dark red color with a light red grid pattern. In the center, there is a large, faint, light red caduceus symbol. To the left and right of the caduceus, there are faint, light red ECG (heart rate) lines. The overall design is clean and professional, typical of a medical journal or publication cover.

AD Publications

Sector-3, MP Colony, Bikaner, Rajasthan, INDIA

www.adpublications.org, www.imjhealth.org, info@imjhealth.org